

Introduction to Quasiconformal Maps  
and Teichmüller Spaces  
Lecture Notes

Note taker: Jacky567

Qiuzhen College, Tsinghua University  
2025 Autumn

# Contents

|          |   |           |
|----------|---|-----------|
| <b>1</b> | <b>Quasiconformal maps</b>                    | <b>1</b>  |
| 1.1      | Differentiable quasiconformal maps            | 1         |
| 1.2      | Extremal length                               | 2         |
| 1.3      | Quadrilateral                                 | 4         |
| 1.4      | Quasiconformal maps                           | 5         |
| 1.5      | Holder continuity                             | 6         |
| 1.6      | quasiconformal maps on Riemann surface        | 15        |
| 1.7      | Topological definition of quasiconformal maps | 15        |
| 1.8      | Analytical properties of quasiconformal maps  | 17        |
| <b>2</b> | <b>Boundary Correspondence</b>                | <b>19</b> |
| 2.1      | Quasi-isometry maps                           | 19        |
| 2.2      | $M$ -condition                                | 21        |
| <b>3</b> | <b>Beltrami differential</b>                  | <b>26</b> |
| 3.1      | Beltrami differential                         | 26        |
| 3.2      | Quasiconformal groups                         | 27        |
| 3.3      | Holomorphic motions                           | 29        |
| 3.4      | Two integral operators                        | 31        |
| 3.5      | Beltrami equations                            | 32        |
| 3.6      | Decomposition of quasiconformal maps          | 34        |
| 3.7      | Dependence on parameter                       | 35        |
| <b>4</b> | <b>Teichmüller space</b>                      | <b>36</b> |
| 4.1      | Integrable holomorphic quadratic differential | 36        |
| 4.2      | Bergman projection                            | 39        |
| 4.3      | Teichmüller spaces                            | 41        |
| 4.4      | Douady-Earle extension                        | 44        |
| 4.5      | Teichmüller space of torus                    | 45        |
| 4.6      | Teichmüller theorem                           | 47        |
| 4.7      | Schwartzian derivative                        | 50        |
| <b>5</b> | <b>Additional topics</b>                      | <b>53</b> |
| 5.1      | Bicanonical embedding                         | 53        |
| 5.2      | Automorphism of $T_g$                         | 54        |
| 5.3      | Quasi-Fuchsian group                          | 55        |
| 5.4      | Complex dynamics                              | 56        |
| <b>A</b> | <b>Quasi-isometry and Mostow rigidity</b>     | <b>58</b> |
| A.1      | Boundary extension of quasi-isometry          | 58        |
| A.2      | Mostow rigidity theorem                       | 62        |

# Chapter 1

## Quasiconformal maps

### 1.1 Differentiable quasiconformal maps

**Def 1.1.1.** Let  $(M_1, d_1), (M_2, d_2)$  be two Riemannian manifold and  $f : M_1 \rightarrow M_2$  is a homeomorphism, define the dilatation

$$D_f : M_1 \rightarrow \mathbb{R}, x \mapsto \overline{\lim}_{r \rightarrow 0} \frac{\beta(f(B_r(x)))}{\alpha(f(B_r(x)))}$$

where  $\alpha(f(B_r(x)))$  is the diameter of the largest ball that can be inscribed in  $f(B_r(x))$  and  $\beta(f(B_r(x)))$  is the diameter of smallest ball circumscribe  $f(B_r(x))$ .

**Def 1.1.2.** We say that  $f$  is  $k$ -quasiconformal if  $D_f(x) \leq k$  for any  $x \in M_1$ .

**Exam 1.1.1.** (1)  $M_1 = M_2 = \mathbb{R}$ , then  $D_f(x) = 1$  for every  $x \in \mathbb{R}$ .

(2)  $M_1 = M_2 = \mathbb{R}^n$  and  $f$  is bi-Lipschitz map, then  $\beta(f(B_r)) \leq Lr, \alpha(f(B_r)) \geq \frac{r}{L}$ , i.e.  $f$  is  $L^2$ -quasiconformal.

(3)  $M_1 = M_2 = \mathbb{C}$  and  $f(re^{i\theta}) = r^2 e^{i\theta}$ , then  $f$  is 2-quasiconformal but not bi-Lipschitz on every neighborhood of 0.

**Prop 1.1.1.** Suppose  $U \subset \mathbb{R}^k$  is a domain and  $f : U \rightarrow f(U)$  is  $C^1$ -diffeomorphism, then the dilatation can given by

$$D_f(p) = \frac{\sup_{\|v\|=1} \|df_p(v)\|}{\inf_{\|v\|=1} \|df_p(v)\|}$$

*Proof.* Since around  $p$ ,

$$f(q) = f(p) + df_p(q - p) + o(\|q - p\|).$$

So for sufficiently small  $r$ ,  $f(B_r(p))$  is closed to the ellipsoid

$$E = f(p) + df_p(B_r(0)) \subset T_{f(p)}f(U).$$

And notice that  $\alpha(E) = 2r \sup_{\|v\|=1} \|df_p(v)\|, \beta(E) = 2r \sup_{\|v\|=1} \|df_p(v)\|$ .

Hence

$$D_f(p) = \overline{\lim}_{r \rightarrow 0} \frac{\beta(f(B_r(x)))}{\alpha(f(B_r(x)))} = \frac{\sup_{\|v\|=1} \|df_p(v)\|}{\inf_{\|v\|=1} \|df_p(v)\|}$$

□

**Coro 1.1.1.** Suppose  $U \subset \mathbb{R}^k$  is a domain and  $f : U \rightarrow f(U)$  is  $C^1$ -diffeomorphism, then on every compact subset of  $U$ ,  $D_f$  is continuous and  $f$  is quasiconformal.

**Prop 1.1.2.** Let  $f : \mathbb{C} \rightarrow \mathbb{C}$ , then


$$Df(z_0) = \frac{|f_z| + |f_{\bar{z}}|}{|f_z| - |f_{\bar{z}}|}(z_0).$$

*Proof.* Since  $df = f_z dz + f_{\bar{z}} d\bar{z}$ .

So we have

$$(|f_z| - |f_{\bar{z}}|)|dz| \leq |dw| \leq (|f_z| + |f_{\bar{z}}|)|dz|.$$

Geometrically,  $df$  maps the unit circle to an ellipse:



and two axes of the ellipse are  $|f_z| + |f_{\bar{z}}|, |f_z| - |f_{\bar{z}}|$ .

Hence we have

$$Df(z) = \frac{|f_z| + |f_{\bar{z}}|}{|f_z| - |f_{\bar{z}}|}$$

□

## 1.2 Extremal length

**Def 1.2.1.** A function  $\rho : \mathbb{C} \rightarrow \mathbb{R}$  is called allowable if

- (1)  $\rho \geq 0$  and measurable,
- (2)  $A(\rho) = \int_{\mathbb{C}} \rho^2(z) dx dy \neq 0, \infty$ .

Let  $\Gamma$  be a family of curves, each  $\gamma \in \Gamma$  is a countable union of open arcs which are rectifiable, define

$$L_{\gamma}(\rho) = \int_{\gamma} \rho(z) |dz|, L(\rho) = \inf_{\gamma \in \Gamma} L_{\gamma}(\rho).$$

The external length of  $\Gamma$  is defined as

$$\lambda(\Gamma) = \sup_{\rho} \frac{L^2(\rho)}{A(\rho)}$$

**Def 1.2.2.** We say  $\Gamma_1 < \Gamma_2$  if every  $\gamma_2$  contains  $\gamma_1$ .

**Exam 1.2.1.** If  $\Gamma_1 \subset \Gamma_2$ , then  $\Gamma_2 < \Gamma_1$ .

**Prop 1.2.1.** (1) If  $\Gamma_1 < \Gamma_2$ , then  $\lambda(\Gamma_1) \leq \lambda(\Gamma_2)$ .

(2) Let  $\Gamma_1 + \Gamma_2 = \{\gamma_1 + \gamma_2 | \gamma_i \in \Gamma_i\}$ , then  $\lambda(\Gamma_1 + \Gamma_2) \geq \lambda(\Gamma_1) + \lambda(\Gamma_2)$ .

(3) If  $\Gamma_1 \cap \Gamma_2 = \emptyset$ , then  $\lambda(\Gamma_1 \cup \Gamma_2)^{-1} \geq \lambda(\Gamma_1)^{-1} + \lambda(\Gamma_2)^{-1}$ .

*Proof.* (1) If  $\gamma_1 \subset \gamma_2$ , then  $L_{\gamma_1}(\rho) \leq L_{\gamma_2}(\rho)$ .

So  $\lambda(\Gamma_1) \leq \lambda(\Gamma_2)$ .

(2) WLOG, assume  $L_i(\rho_i) = A(\rho_i)$ .

Let  $\rho = \max(\rho_1, \rho_2)$ , then

$$\begin{aligned} L(\rho) &\geq L_1(\rho_1) + L_2(\rho_2) = A(\rho_1) + A(\rho_2) \\ A(\rho) &\leq A(\rho_1) + A(\rho_2) \end{aligned}$$

So

$$\lambda(\Gamma_1 + \Gamma_2) = \sup_{\rho} \frac{L^1(\rho)}{A(\rho)} \geq A(\rho_1) + A(\rho_2) = \lambda(\Gamma_1) + \lambda(\Gamma_2).$$

(3) Let  $E_1, E_2$  be two complementary measurable sets with  $\Gamma_i \subset E_i$ .

For allowable  $\rho$ , take  $\rho_i = \rho \cdot \chi_{E_i}$ .

Then  $L_1(\rho_1) \geq L(\rho), L_2(\rho_2) \geq L(\rho)$  and  $A(\rho) = A(\rho_1) + A(\rho_2)$ .

So

$$\frac{A(\rho)}{L^2(\rho)} \geq \frac{A(\rho_1)}{L_1^2(\rho_1)} + \frac{A(\rho_2)}{L_2^2(\rho_2)}$$

Hence  $\lambda(\Gamma_1 \cup \Gamma_2)^{-1} \geq \lambda(\Gamma_1)^{-1} + \lambda(\Gamma_2)^{-1}$

□

**Exam 1.2.2.** Let  $\Gamma$  be the set of all arcs in an annulus  $r_1 \leq |z| \leq r_2$  which join the boundary circles, then

$$\int_0^{2\pi} \int_{r_1}^{r_2} \rho(re^{i\theta}) dr d\theta \geq \int_0^{2\pi} L(\rho) d\theta = 2\pi L(\rho).$$

And by Cauchy inequality,

$$4\pi^2 L^2(\rho) \leq \int_0^{2\pi} \int_{r_1}^{r_2} \frac{1}{r} dr d\theta \cdot \int_0^{2\pi} \int_{r_1}^{r_2} \rho^2 r dr d\theta \leq 2\pi \log \frac{r_2}{r_1} A(\rho)$$

Hence  $\lambda(\Gamma) = \frac{1}{2\pi} \log \frac{r_2}{r_1}$ .

**Thm 1.2.1.** Suppose  $\Gamma \subset U \subset \mathbb{R}^2, f : U \rightarrow U'$  is diffeomorphism and  $\Gamma' = f(\Gamma)$ , if  $f$  is  $k$ -quasiconformal, then  $\frac{\lambda(\Gamma)}{k} \leq \lambda(\Gamma') \leq k\lambda(\Gamma)$ .

*Proof.* Given density  $\rho$  on  $U_1$ , we define

$$\rho'(\zeta) = \left( \frac{\rho}{|f_z| - |f_{\bar{z}}|} \right) (z)$$

with  $f(z) = \zeta$ , then

$$|d\zeta| \geq (|f_z| - |f_{\bar{z}}|) |dz|.$$

So

$$\int_{\gamma'} \rho' |d\zeta| \geq \int_{\gamma} \rho |dz|.$$

Thus  $L_{\gamma'}(\rho') \geq L_{\gamma}(\rho)$ , i.e.  $L(\rho') \geq L(\rho)$ .

On the other hand,

$$A(\rho') = \int_{U'} \rho' d\zeta_1 d\zeta_2 = \int_U (\rho' \circ f)^2 J_f dx dy = \int_U \rho^2 \frac{|f_z| + |f_{\bar{z}}|}{|f_z| - |f_{\bar{z}}|} dx dy \leq kA(\rho)$$

Therefore  $\lambda(\Gamma') \geq \frac{\lambda(\Gamma)}{k}$ .

And consider

$$\rho''(\zeta) = \left( \frac{\rho}{|f_z| + |f_{\bar{z}}|} \right) (z),$$


we have  $\lambda(\Gamma') \leq k\lambda(\Gamma)$ .

□

**Coro 1.2.1.**  $\lambda(\Gamma)$  is a conformal invariant.

### 1.3 Quadrilateral

**Def 1.3.1.**  $Q$ -quadrilateral is a Jordan domain with a pair of disjoint closed arcs on the boundary (the b-sides).



**Lemma 1.3.1.** *There exists  $M > 0$  and a conformal map  $f : Q \rightarrow R$ , where  $R$  is a rectangle, such that the b-sides are mapped to the vertical sides.*

*Proof.* follows from Riemann mapping theorem and Schwarz-Christoffel formula. □

*Remark 1.3.1.* In this note, we assume the width of  $R$  is 1 and the length is  $M$ .

**Prop 1.3.1.** *If  $f : R \rightarrow R'$  is conformal and mapping corner to corner, then the length  $M = M'$  and  $f$  is identity.*

*Proof.* By reflecting, we can extend  $f$  to whole  $\mathbb{C}$ .

And since

$$\lim_{|z| \rightarrow \infty} \frac{|f(z)|}{|z|} < +\infty$$

So  $f$  must be a degree-1 polynomial.

Notice that  $f(0) = 0, f(M) = M', f(i) = i$ .

Hence  $f$  must be identity and  $M = M'$ . □

**Def 1.3.2.** The modular of a rectangle  $R$  is  $M$ , denoted by  $m(R) = M$ .

More generally, given  $Q$ , we define  $m(Q) = m(R)$  when we have conformal map  $f : Q \rightarrow R$ .

**Prop 1.3.2.**  $\lambda(\Gamma) = m(Q)$ , where  $\Gamma$  is the family of arcs connecting the b-sides of  $Q$ .

*Proof.* By corollary 1.2.1, we only need to prove this for  $Q = R$  is a rectangle, so

$$\int_0^1 \int_0^M \rho(x + iy) dx dy \geq \int_0^1 L(\rho) dy = L(\rho).$$

And by Cauchy inequality,

$$L^2(\rho) \leq \int_0^1 \int_0^M dx dy \cdot \int_0^1 \int_0^M \rho^2 dx dy \leq MA(\rho).$$

Hence  $\lambda(\Gamma) = M = m(Q)$ . □

**Def 1.3.3.** Let  $\Gamma$  be the family of arcs connecting the b-side, define

$$s_b = \inf_{\gamma \in \Gamma} L_\gamma(\rho)$$

where  $\rho$  is the Euclidean density, and similarly, we can define  $s_a$ .

**Thm 1.3.1** (Rengel inequality).

$$\frac{s_b^2(Q)}{A(Q)} \leq m(Q) \leq \frac{A(Q)}{s_a^2(Q)}.$$

*The equality holds iff  $Q$  is a rectangle.*

*Proof.* Let  $\Gamma_b$  be the family of arcs connecting the b-sides of  $Q$ , then

$$\frac{s_b^2(Q)}{A(Q)} \leq \lambda(\Gamma_b) = m(Q).$$

On the other hand, we consider quadrilateral  $Q'$  whose b-sides are the a-sides of  $Q$ , then

$$\frac{s_a^2(Q)}{A(Q)} = \frac{s_b^2(Q')}{A(Q')} = m(R') = \frac{1}{m(R)}.$$

Hence we have

$$\frac{s_b^2(Q)}{A(Q)} \leq m(Q) \leq \frac{A(Q)}{s_a^2(Q)}.$$

□

## 1.4 Quasiconformal maps

**Def 1.4.1.** Let  $\Omega \subset \mathbb{C}$  be a domain, and suppose  $f : \Omega \rightarrow f(\Omega)$  is a homeomorphism, we say that  $f$  is  $k$ -quasiconformal if

$$\frac{m(Q)}{k} \leq m(f(Q)) \leq km(Q)$$

for every quadrilateral  $Q \subset \Omega$ .

**Prop 1.4.1.** (1) If  $f$  is  $k$ -quasiconformal,  $f^{-1}$  is  $k$ -quasiconformal

(2) If  $f_1$  is  $k_1$ -quasiconformal and  $f_2$  is  $k_2$ -quasiconformal, then  $f_1 \circ f_2$  is  $k_1k_2$ -quasiconformal.

(3) Conformal map is 1-quasiconformal.

*Proof.* (1)

$$\frac{m(f^{-1}(Q))}{k} \leq m(Q) \leq km(f^{-1}(Q)).$$

(2)

$$\frac{m(Q)}{k_1k_2} \leq \frac{m(f_2(Q))}{k_1} \leq m(f_1 \circ f_2(Q)) \leq k_1m(f_2(Q)) \leq k_1k_2m(Q).$$

(3) Let  $R$  be a rectangle and WLOG, assume  $f(R) = R'$  is also an rectangle otherwise we can composite a conformal map.

Then by proposition 1.3.1,  $f$  is identity, *i.e.*  $f$  is 1-quasiconformal.

□

**Thm 1.4.1.** Every 1-quasiconformal map is conformal.

*Proof.* Let  $R$  be a rectangle and WLOG, assume  $f(R) = R'$  is also an rectangle otherwise we can composite a conformal map.

Consider a vertical line  $l$  that divides  $R$  into  $Q_1$  and  $Q_2$  and let  $Q'_i = f(Q_i)$ .

Then  $m(Q'_i) = m(Q_i)$  and  $f(l)$  is an arc connecting the a-sides.

By Rengel inequality,

$$m(Q'_i) \leq \frac{A(Q'_i)}{s_a^2(Q'_i)} = A(Q'_i).$$

Suppose  $f(l)$  is not a vertical line, *i.e.*  $Q_1, Q_2$  are not rectangle.

Then  $m(Q'_1) + m(Q'_2) < A(Q'_1) + A(Q'_2) = A(R') = A(R)$ , contradiction!

So  $f$  is identity.

For general cases,  $f$  must be conformal.

□

**Prop 1.4.2.** *If  $f : \Omega \rightarrow f(\Omega)$  is  $C^1$ , then  $f$  is  $k$ -quasiconformal in the  $C^1$  sense iff  $f$  is  $k$ -quasiconformal in geometric sense.*

*Proof.*  $\Rightarrow$ : Let  $Q$  be a quadrilateral in  $\Omega$ .

WLOG, we assume  $R = Q$  and  $R' = f(Q)$  are rectangles, then

$$|f_x|^2 \leq (|f_z| + |f_{\bar{z}}|)^2 \leq kJ_f$$

So

$$\begin{aligned} m(R') = M' &= \int_R J_f(z) dx dy \\ &\geq \frac{1}{k} \int_R |f_x|^2(z) dx dy \\ &= \frac{1}{Mk} \int_0^1 \left( \int_0^M dx \cdot \int_0^M |f_x|^2(z) dx \right) dy \\ &\geq \frac{1}{Mk} \int_0^1 \left( \int_0^M |f_x|(x) dx \right)^2 dy \geq \frac{1}{Mk} (M')^2 \end{aligned}$$

Hence  $M' \leq kM$ , *i.e.*  $f$  is  $k$ -quasiconformal in geometric sense.

$\Leftarrow$ : We first assume  $f(z) = |f_z(0)|z + |f_{\bar{z}}(0)|\bar{z} + o(z)$ .

Consider square  $R_\delta$  with length  $\delta$  and  $R'_\delta = f(R_\delta)$ .

Then

$$\begin{aligned} s_b(R'_\delta) &= \delta(|f_z(0)| + |f_{\bar{z}}(0)|) + o(\delta) \\ A(R'_\delta) &= \delta^2(|f_z(0)|^2 - |f_{\bar{z}}(0)|^2) + o(\delta^2). \end{aligned}$$

So by Rengel inequality,

$$m(R'_\delta) \geq \frac{s_b^2(R'_\delta)}{A(R'_\delta)} = \frac{\delta^2 (|f_z(0)| + |f_{\bar{z}}(0)|)^2 + o(\delta^2)}{\delta^2 (|f_z(0)| - |f_{\bar{z}}(0)|)^2 + o(\delta^2)} = D_f(0) + O(1).$$

Hence  $D_f(0) \leq k$ , *i.e.*  $f$  is  $k$ -quasiconformal in  $C^1$ -sense.

For the general case, Let  $\arg f_z(0) = \theta$ ,  $\arg f_{\bar{z}}(0) = \varphi$  and

$$g(z) = e^{-i\frac{\theta+\varphi}{2}} f \left( e^{i\frac{\varphi-\theta}{2}} z \right).$$

Then

$$\begin{aligned} g_z(0) &= e^{-i\frac{\theta+\varphi}{2}} f_z(0) \cdot e^{i\frac{\varphi-\theta}{2}} = |f_z(0)| \\ g_{\bar{z}}(0) &= e^{-i\frac{\theta+\varphi}{2}} f_{\bar{z}}(0) \cdot \overline{e^{i\frac{\varphi-\theta}{2}}} = |f_{\bar{z}}(0)| \end{aligned}$$

And since the rotations are conformal, *i.e.* 1-quasiconformal.

Hence  $f$  is  $k$ -quasiconformal in  $C^1$ -sense as  $g$  is. □

## 1.5 Holder continuity

**Def 1.5.1.** Let  $\Omega$  be a topological annulus (doubly connected region) and  $C_1, C_2$  be the bounded, unbounded region of the component resp.

We say the closed curve  $\gamma$  in  $G$  separates  $C_1$  and  $C_2$  if  $\gamma$  has non-zero winding number about the points of  $C_1$ .

We denote  $\Gamma = \{\gamma | \gamma \text{ separates } C_1 \text{ and } C_2\}$  and define the modular  $m(\Omega) = \lambda(\Gamma)^{-1}$ .

**Prop 1.5.1.** Let  $\Omega$  be a topological annulus and  $f$  maps  $\Omega$  conformally to annulus  $\{r_1 < |z| < r_2\}$ , then

$$m(\Omega) = \inf_{\rho} \frac{A(\rho)}{L^2(\rho)} = \frac{1}{2\pi} \log \frac{r_2}{r_1}$$

and reach the maximum at

$$\rho_0 = \left| \frac{f'}{f} \right|.$$

*Proof.* For any allowable  $\rho$ , take  $\rho_1(f(z)) = \rho(z) \cdot \rho_0(z)^{-1}$ .

For  $\gamma_r \subset \Omega$  such that  $f(\gamma_r) = \{|z| = r\}$ , we have


$$\begin{aligned} L_{\gamma_r}(\rho) &= \int_{\gamma_r} \rho |dz| = \int_{|z|=r} \rho_1 \frac{1}{r} |dz| = \int_0^{2\pi} \rho_1(r e^{i\theta}) d\theta. \\ \iint_{\Omega} \rho^2 dx dy &= \int_{r_1}^{r_2} \int_0^{2\pi} \rho_1^2(r e^{i\theta}) \frac{1}{r} d\theta dr \\ &\geq \frac{1}{2\pi} \int_{r_1}^{r_2} \frac{1}{r} \left( \int_0^{2\pi} \rho_1(r e^{i\theta}) d\theta \right)^2 dr \\ &\geq \frac{1}{2\pi} \int_{r_1}^{r_2} \frac{L^2(\rho)}{r} dr = \frac{L^2(\rho)}{2\pi} \log \frac{r_2}{r_1} \end{aligned}$$

Hence we obtain the conclusion. □

**Coro 1.5.1.** Consider  $\Omega_r = \{|z| \leq 1\} \setminus [0, r]$ , prove that

$$m(\Omega_r) = \frac{\mu(r)}{2\pi} = \frac{1}{4} \frac{I(\sqrt{1-r^2})}{I(r)}, I(r) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-r^2x^2)}}.$$

*Proof.*



Consider Möbius transformation

$$f(z) = \frac{z - \sqrt{a}}{-\sqrt{a}z + 1}, \frac{2\sqrt{a}}{a+1} = r$$

Then  $f(\Omega_r) = \{|z| \leq 1\} \setminus [-\sqrt{a}, \sqrt{a}]$ .

Let  $\Omega^+ = f(\Omega_r) \cap \{\text{Im } z \geq 0\}$  and consider

$$g(z) = \int_0^{\frac{z}{\sqrt{a}}} \frac{dt}{\sqrt{(1-t^2)(1-a^2t^2)}}.$$

We can verify that  $g(\Omega_r^+) = \{-K \leq \text{Re } z \leq K, 0 < \text{Im } z < K'\}$  is a rectangle with

$$K = I(a), K' = \frac{1}{2} I(\sqrt{1-a^2}).$$

And  $g((\sqrt{a}, 1)) = (K, K + iK'), g(-1, -\sqrt{a}) = (-K, -K + iK')$ .

Consider

$$h(z) = te^{\frac{i\pi}{2K}(K-z)}.$$

Then  $h(g(\Omega_r^+)) = \{t < |z| < 1, \text{Im } z \geq 0\}$  with

$$t = e^{-\frac{K'\pi}{2K}}.$$

Notice that  $h \circ g$  maps  $(-1, -\sqrt{a}) \cup (\sqrt{a}, 1)$  to  $(-1, -t) \cup (t, 1)$ .

So we obtain a conformal map  $g_0$  by reflect  $h \circ g$  along real axis, which maps  $f(\Omega_r)$  to annulus  $\{t < |z| < 1\}$ .

By proposition 1.5.1,

$$m(\Omega_r) = -\frac{1}{2\pi} \log t = \frac{1}{8} \frac{I(\sqrt{1-a^2})}{I(a)}.$$

And since  $I(r) = (1+a)I(a), I(\sqrt{1-r^2}) = I\left(\frac{1-a}{1+a}\right) = \frac{1}{2}(1+a)I(\sqrt{1-a^2})$ .

Hence

$$m(\Omega_r) = \frac{1}{8} \frac{I(\sqrt{1-a^2})}{I(a)} = \frac{1}{4} \frac{I(\sqrt{1-r^2})}{I(r)}.$$

□


*Remark 1.5.1.* We call  $\Omega_r$  the Grötzsch extremal region.

By the prove of the above corollary, we have

$$\frac{\mu(r)}{2} = \mu\left(\frac{2\sqrt{r}}{1+r}\right).$$

More details about the properties of elliptic functions can be found in *Conformal Invariants, Inequalities, and Quasiconformal Maps*.

**Prop 1.5.2.** Let  $\Omega = \{|z| \leq 1\} \setminus c$  where  $c$  is an arc connecting 0 and  $r$ , then  $m(\Omega) \leq m(\Omega_r)$ .



*Proof.* By corollary 1.5.1, we can consider the conformal maps  $f : \overline{\Omega_r^+} \rightarrow \{t \leq |z| \leq 1, \text{Im } z \geq 0\}$  and the allowable function  $\rho_0$ .

Let  $\gamma$  be an arbitrary curve in  $\Gamma$  and divides it into two arcs  $\gamma_1$  and  $\gamma_2$  which both have one endpoint on each of the segments  $(-1, 0)$  and  $(r, 1)$ .

Let  $\gamma_i^+$  be obtained by reflecting the part of  $\gamma_i$  below the real axis.

Since it is well-defined to extend  $f$  on upper half disc.

So  $f(\gamma_i^+)$  is an arc which has one endpoint on each of the segments  $(-1, -t)$  and  $(t, 1)$ .

Notice that  $\rho$  is symmetric w.r.t. real axis.

So  $L_{\gamma_i}(\rho_0) = L_{\gamma_i^+}(\rho_0) \geq \pi$ .

Hence

$$\lambda(\Gamma) \geq \frac{\inf_{\gamma \in \Gamma} L_\gamma^2(\rho_0)}{-2\pi \log t} \geq \frac{(2\pi)^2}{-2\pi \log t} = \lambda(\Gamma_r)$$

□

**Prop 1.5.3.**  $d(0, w) = \log \left( \frac{1+|w|}{1-|w|} \right)$  for  $w \in \mathbb{D}$ .

**Prop 1.5.4.** Suppose  $f : \mathbb{D} \rightarrow \mathbb{D}$  and  $f$  is 2-quasiconformal, then

$$\frac{1}{2}d(z, w) - \log 2 \leq d(f(z), f(w)) \leq 2d(z, w) + 2 \log 2.$$

*Proof.* WLOG, assume  $w = f(w) = 0$ .

Since  $m(\Omega) \geq \frac{1}{2}m(\Omega_2)$ , we have

$$\frac{\mu(|z|)}{2} \leq \mu(|f(z)|).$$

And so

$$|f(z)| \leq \mu^{-1} \left( \frac{\mu(|z|)}{2} \right) = \frac{2\sqrt{|z|}}{1+|z|}.$$

Hence

$$d(0, |f(z)|) \leq d \left( 0, \frac{2\sqrt{|z|}}{1+|z|} \right) = 2d(0, \sqrt{z}) \leq 2d(0, z) + 2 \log 2.$$

And the other side of inequality follows from 2-quasiconformal map  $f^{-1}$ . □

**Prop 1.5.5.** Suppose  $f : \mathbb{D} \rightarrow \mathbb{D}$  and  $f$  is  $2^n$ -quasiconformal, then

$$\frac{1}{2^n}d(z, w) - (2 - 2^{1-n}) \log 2 \leq d(f(z), f(w)) \leq 2^n d(z, w) + (2^{n+1} - 2) \log 2.$$

*Proof.* Let

$$\phi(z) = \mu^{-1} \left( \frac{\mu(|z|)}{2} \right) = \frac{2\sqrt{|z|}}{1+|z|}$$

Similar to proposition 1.5.4, we have

$$|f(z)| \leq \mu^{-1} \left( \frac{\mu(|z|)}{2^n} \right) = \phi^{(n)}(z),$$

which is the  $n$ -th iterate of the function  $\phi$ .

Then

$$\begin{aligned} d(0, |f(z)|) &\leq d(0, \phi^{(n)}(z)) \\ &\leq 2d(0, \phi^{(n-1)}(z)) + 2 \log 2 \\ &\leq \dots \\ &\leq 2^n d(0, z) + (2^{n+1} - 2) \log 2 \end{aligned}$$

And the other side of inequality follows from  $2^n$ -quasiconformal map  $f^{-1}$ . □

**Lemma 1.5.1.** Given  $k$  and a point  $p$  on a hyperbolic geodesic  $\gamma$ , there is an  $L$ -quasiconformal diffeomorphism  $f : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  which fixes  $p$  and sends  $\gamma$  to itself, stretching hyperbolic distances along  $\gamma$  by a factor  $k$  and stretching all other distances by a factor between 1 and  $k$ .

*Proof.* WLOG, assume  $p = i$  and  $\gamma$  is the image axis, then take

$$f(re^{i\theta}) = r^k e^{i\theta}.$$

Then

$$\begin{aligned} |fz| &= \frac{k+1}{2}|z|^{k-1}, |f\bar{z}| = \frac{k-1}{2}|z|^{k-1} \\ D_f(z) &= \frac{\frac{k+1}{2}|z|^{k-1} + \frac{k-1}{2}|z|^{k-1}}{\frac{k+1}{2}|z|^{k-1} - \frac{k-1}{2}|z|^{k-1}} = k \end{aligned}$$

□

**Thm 1.5.1.** *Suppose  $f : \mathbb{D} \rightarrow \mathbb{D}$  and  $f$  is  $k$ -quasiconformal, then*

$$\frac{1}{k}d(z, w) - A \leq d(f(z), f(w)) \leq kd(z, w) + A,$$

where  $A = A(k)$  only depends on  $k$ .

*Proof.* WLOG, assume  $w = f(w) = 0$  and let  $2^{n-1} < k \leq 2^n$ .

Then by lemma 1.5.1, there exists a  $\frac{2^n}{k}$ -quasiconformal  $g$  that fix 0 and maps the geodesic connecting 0 and  $f(z)$  to itself.

So by proposition 1.4.1,  $g \circ f$  is  $2^n$ -quasiconformal, i.e. by proposition 1.5.5,

$$\frac{2^n}{k}d(f(z), 0) = d(g \circ f(z), 0) \leq 2^n d(z, 0) + (2^{n+1} - 2) \log 2.$$

Therefore

$$d(0, f(z)) \leq kd(0, z) + k(2 - 2^{1-n}) \log 2.$$

And the other side of inequality follows from  $k$ -quasiconformal map  $f^{-1}$ .

□

**Lemma 1.5.2.** *Consider topological annulus*

$$\Omega'_p = \mathbb{C} \setminus ([-1, 0] \cup [p, +\infty)),$$

let  $\Omega$  be a topological annulus and  $C_1, C_2$  be the two components of its complement, if

$$\{-1, 0\} \subset C_1, \{p, \infty\} \subset C_2,$$

then

$$m(\Omega) \leq m(\Omega'_p) = \frac{1}{2\pi} \mu \left( 2p + 1 - 2\sqrt{p^2 + p} \right).$$

*Proof.* Let  $f : \mathbb{D}^2 \rightarrow C_1 \cup \Omega$  be a conformal map with  $f(0) = 0$  and  $f(a) = -1$ .

Then by Koebe quarter theorem,  $|f'(0)| = 4p$ .

And by Koebe distortion theorem,

$$1 = |f(a)| \leq \frac{|a||f'(0)|}{(1-|a|)^2} \leq \frac{4p|a|}{(1-|a|)^2}.$$

So  $|a| \geq |a_0|$  where  $a_0$  belongs to the case  $\Omega = \Omega'_p$ .

by proposition 1.5.2,  $m(\Omega) = m(f^{-1}(\Omega)) \leq m(\Omega_a) \leq m(\Omega_{a_0}) = m(\Omega'_p)$ .

Moreover, we have

$$a_0 = 2p + 1 - 2\sqrt{p^2 + p}, m(\Omega'_p) = \frac{1}{2\pi} \mu \left( 2p + 1 - 2\sqrt{p^2 + p} \right).$$

□

*Remark 1.5.2.*  $\Omega_p$  is called the Teichmüller extremal region.

Let  $\Omega_r^-$  be the reflection of  $\Omega_r$  along  $\{|z| = 1\}$  and  $\Omega = \Omega_r \cup \Omega_r^- \cup \{|z| = 1\}$ .

Then  $f(\Omega) = \Omega'_{r^{-2}-1}$  with  $f(z) = \frac{z}{r} - 1$  and so

$$2\mu(r) = \mu\left(2r^{-2} - 1 - 2\sqrt{r^{-4} - r^{-2}}\right).$$

$$2\mu\left(\frac{2\sqrt{a}}{a+1}\right) = \mu(a).$$

We can actually use this conclusion in corollary 1.5.1 without creating circular argument.

**Lemma 1.5.3.** *Consider topological annulus*

$$\Omega'_\lambda = \mathbb{C} \setminus \left( \left\{ |z| = 1, |\arg z - \pi| \leq \arcsin \frac{\lambda}{2} \right\} \cup [0, +\infty) \right),$$

let  $\Omega$  be a topological annulus and  $C_1, C_2$  be the two components of its complement, if

$$\text{diam}(C_1 \cap \{|z| \leq 1\}) \geq \lambda, \{0, \infty\} \subset C_2,$$

then

$$m(\Omega) \leq m(\Omega'_\lambda) = \frac{1}{2\pi} \mu\left(\frac{\sqrt{4+2\lambda} - \sqrt{4-2\lambda}}{4}\right).$$

*Proof.* Consider the map  $f(z) = \sqrt{z}$ .

On the image plane, we get a figure which is symmetric w.r.t. the origin with two component images of  $C_1$  and two of  $C_2$ , denote them by  $C_1^+, C_1^-, C_2^+, C_2^-$ .

Take  $\Omega' = \mathbb{C} \setminus (C_1^+ \cup C_1^-)$ .

Then  $m(\Omega) \leq \frac{1}{2}m(\Omega')$ .

And since there exists  $z_1, z_2$  such that  $|z_1|, |z_2| \leq 1, |z_1 - z_2| \geq \lambda$ .

Let  $w_1, w_2 \in C_1^+$  are the image and

$$g(z) = \frac{z + w_1}{z - w_1} \frac{w_1 + w_2}{w_1 - w_2}.$$

Then

$$g(-w_1) = 0, g(-w_2) = 1, g(w_1) = \infty, g(w_2) = -u^2 = -\left(\frac{w_1 + w_2}{w_1 - w_2}\right)^2.$$

Notice that

$$|z_1 + z_2|^2 = 2(|z_1|^2 + |z_2|^2) - |z_1 - z_2|^2 \leq 4 - \lambda^2.$$

So we can obtain

$$|u| - \frac{1}{u} = \frac{2(z_1 + z_2)}{z_2 - z_1} \leq \frac{2}{\lambda} \sqrt{4 - \lambda^2}, |u| \leq \frac{2 + \sqrt{4 - \lambda^2}}{\lambda},$$

Hence by lemma 1.5.2,

$$m(\Omega) \leq \frac{1}{2}m(\Omega') \leq \frac{1}{2}m(\Omega'_{|u|^2}) = \frac{1}{2\pi} \mu\left(\sqrt{\frac{1}{|u|^2 + 1}}\right) \leq \frac{1}{2\pi} \mu\left(\frac{\sqrt{4+2\lambda} - \sqrt{4-2\lambda}}{4}\right).$$

And the equality holds when  $\Omega = \Omega'_\lambda$ , i.e.  $m(\Omega) \leq m(\Omega'_\lambda)$  □

**Lemma 1.5.4.**

$$2 \log \frac{1 + \sqrt{1 - r^2}}{r} \leq \mu(r) \leq \log \frac{4}{r}.$$

*Proof.* By Möbius transformation

$$f(z) = \frac{z - a}{a - a^2z}, \frac{2a}{a^2 + 1} = r,$$

we map  $\Omega_r$  to  $\Omega = \{|z| \leq a\} \setminus [-1, 1]$ .

And take conformal map

$$g(z) = \frac{1}{2} \left( z + \frac{1}{z} \right).$$

Then it maps annulus  $\{1 < |z| < \rho\}$  to ellipse  $E_\rho$  with axes  $\rho \pm \rho^{-1}$  and slit along  $[-1, 1]$ .

When  $\rho + \rho^{-1} \leq 2a$ ,  $E_\rho \subset \Omega$  and so

$$\frac{\log \rho}{2\pi} = m(E_\rho) \leq m(\Omega) = \frac{\mu(r)}{2\pi}.$$

When  $\rho - \rho^{-1} \geq 2a$ ,  $E_\rho \subset \Omega$  and so

$$\frac{\log \rho}{2\pi} = m(E_\rho) \geq m(\Omega) = \frac{\mu(r)}{2\pi}.$$

Notice that  $\rho = 2a - r, 4r^{-1}$  satisfies the condition of first and second case resp.

So

$$2 \log \frac{1 + \sqrt{1 - r^2}}{r} \leq \mu(r) \leq \log \frac{4}{r}.$$

□

**Thm 1.5.2** (Mori). *Suppose  $f : \mathbb{D} \rightarrow \mathbb{D}$ ,  $f(0) = 0$  and  $f$  is  $k$ -quasiconformal, then*

$$|f(z_1) - f(z_2)| \leq 16|z_1 - z_2|^{\frac{1}{k}}$$

for  $|z_1|, |z_2| \leq 1$ .

*Proof.* We first consider the case that  $f$  can be continuously extended to the closed disk.

And then we can extend  $f$  to a  $k$ -quasiconformal map on  $\mathbb{C}$  by

$$f\left(\frac{1}{\bar{z}}\right) = \frac{1}{\bar{f(z)}}.$$

If  $z_1 - z_2 \geq 1$ , then it is trivial.

So we assume  $z_1 - z_2 < 1$ .

Construct an annulus  $A$  whose inner circle has the segment  $z_1, z_2$  for diameter and whose outer circle is a concentric circle of radius  $\frac{1}{2}$ .

Let  $w_1 = f(z_1), w_2 = f(z_2)$  and  $C_1, C_2$  be the two components of complement of  $A$ .

If  $A \subset \mathbb{D}^2$ , then consider

$$g(z) = \frac{z - w_1}{1 - \bar{w}_1 z}.$$

So  $g \circ f(A)$  is a topological annulus and  $0, g(f(z_2)) \in \partial(g \circ f(A))$  and we can obtain

$$\begin{aligned} \frac{1}{2\pi} \log \frac{1}{|z_2 - z_1|} &= m(A) \leq km(g \circ f(A)) \\ &\leq \frac{k}{2\pi} \mu \left( \left| \frac{w_2 - w_1}{1 - \bar{w}_1 w_2} \right| \right) \\ &\leq \frac{k}{2\pi} \log \frac{8}{|w_2 - w_1|} \end{aligned}$$

If  $A \setminus \mathbb{D}^2 \neq \emptyset$ , then  $0 \notin A$ .

So  $f(A)$  is a topological annulus and  $\{w_1, w_2\} \subset f(C_1) \cap \{|z| \leq 1\}$ ,  $\{0, \infty\} \subset f(C_2)$ .

Therefore by lemma 1.5.3,

$$\begin{aligned} \frac{1}{2\pi} \log \frac{1}{z_2 - z_1} &= m(A) \leq km(f(A)) \\ &\leq \frac{k}{2\pi} \mu \left( \frac{\sqrt{4 + 2|w_1 - w_2|} - \sqrt{4 - 2|w_1 - w_2|}}{4} \right) \\ &\leq \frac{k}{2\pi} \log \frac{4\sqrt{8 + 2\sqrt{16 - 4|w_1 - w_2|}}}{|w_1 - w_2|} \\ &< \frac{k}{2\pi} \log \frac{16}{|w_1 - w_2|}. \end{aligned}$$

Hence we obtain the desire formula.

For the general case, we consider  $\mathbb{D}_r = \{|z| < r\}$  and  $g_r(z) = r^{-1}z$ .

Then  $g_r \circ f(rz)$  is  $K$ -quasiconformal and continuous on  $|z| = 1$ .

By the above case, we have

$$|g_r \circ f(rz_2) - g_r \circ f(rz_1)| < 16|z_2 - z_1|^{\frac{1}{k}}.$$

As  $r \rightarrow 1$ ,  $g_r \rightarrow \text{Id}$ , i.e. we obtain

$$|f(z_2) - f(z_1)| \leq 16|z_2 - z_1|^{\frac{1}{k}}$$

So  $f$  is Hölder continuous, i.e. it can be extended to boundary continuously.  $\square$

**Coro 1.5.2.** Suppose  $f : \mathbb{D} \rightarrow \mathbb{D}$  and  $f$  is  $k$ -quasiconformal, then  $f$  can be extended to a homeomorphism of the closed disk.

*Proof.* By Mori theorem,  $f$  is Hölder continuous.

So it can be continuously extended to the closed disk.

And since continuous bijection from compact set to Hausdorff space is homeomorphism.

Hence  $f : \mathbb{D} \rightarrow \bar{\mathbb{D}}$  is homeomorphism.  $\square$

**Coro 1.5.3.** Suppose  $f_n : \mathbb{D} \rightarrow \mathbb{D}$ ,  $f_n(0) = 0$ , if  $f_n$  are  $k$ -quasiconformal, then  $\{f_n\}$  has a subsequence such that  $f_{i_n} \rightrightarrows f$  and  $f$  is  $k$ -quasiconformal.

*Proof.* By Mori theorem,  $\{f_n\}$  are equicontinuous.

So by Ascoli-Arzela theorem,  $\{f_n\}$  has a subsequence  $f_{i_n} \rightrightarrows f$ .

Similarly, there exists a subsequence  $f_{j_n}^{-1} \rightrightarrows g$ .

WLOG, assume  $i_n = j_n$  otherwise, we can take a subsequence in  $\{i_n\} \cap \{j_n\}$ .

So  $f^{-1} = g$ , i.e.  $f$  is homeomorphic.

For an arbitrary quadrilateral  $Q \subset \mathbb{D}$ , take a sequence of  $Q_n \subset Q$  such that  $f_{i_n}(Q_n) \subset f(Q)$ .

Then  $m(f_{i_n}(Q_n)) \leq km(Q_n)$ .

And since  $f_{i_n} \rightrightarrows f$ , so we have

$$m(f(Q)) = \lim_{n \rightarrow \infty} m(f_{i_n}(Q_n)) \leq \lim_{n \rightarrow \infty} km(Q_n) = km(Q)$$

Hence  $f$  is  $k$ -quasiconformal.  $\square$

**Prop 1.5.6.** Suppose  $f : \bar{\mathbb{D}} \rightarrow \bar{\mathbb{D}}$  is  $k$ -quasiconformal and fixes  $p, q, r \in \mathbb{S}^1$ , then  $d(0, f(0)) \leq C$ , where  $C = C(k)$  only depends on  $k$ .

*Proof.* Suppose  $f_n : \mathbb{D} \rightarrow \mathbb{D}$  is  $k$ -quasiconformal,  $f_n$  fixes  $p, q, r$  and  $d(0, f_n(0)) \rightarrow \infty$ .

WLOG, we assume  $\{f_n(0)\}$  converge to a point on  $\partial\mathbb{D}$  and it is not  $p, q$ .

Let  $A_n \in \text{Isom}(\mathbb{D})$  such that  $A_n(f_n(0)) = 0$  and  $g_n = A_n \circ f_n$ .

Then we obtain

$$|g_n(p) - g_n(q)| = \frac{(1 - |f_n(0)|^2) |p - q|}{\left| \frac{1 - \overline{f_n(0)}p}{1 - \overline{f_n(0)}q} \right|} = \frac{(1 - |f_n(0)|^2) |p - q|}{|p - f_n(0)||q - f_n(0)|} \rightarrow 0.$$

By Mori theorem,  $|p - q| \leq 16|g_n(p) - g_n(q)|^{\frac{1}{k}}$ , contradiction!  $\square$

**Thm 1.5.3.** *Given a compact set  $E \subset \mathbb{C}$  and two point  $p \neq q \in E$ , there exists  $C = C(E)$ , such that if  $f : \mathbb{C} \rightarrow \mathbb{C}$ ,  $f(p) = p$ ,  $f(q) = q$  and  $f$  is  $k$ -quasiconformal, then for  $z_1, z_2 \in E$ ,*

$$|f(z_1) - f(z_2)| \leq C|z_1 - z_2|^{\frac{1}{k}}$$

*Proof.* Take a simply connected region  $\Omega \supset E$ .

Let  $g : \mathbb{D} \rightarrow \Omega$ ,  $h : f(\Omega) \rightarrow \mathbb{D}$  be conformal maps such that  $g(0) = p$ ,  $g(\alpha) = q$ ,  $h(p) = 0$ ,  $h(q) = \beta$  with  $\alpha, \beta \in \mathbb{R}$  and denote  $\phi = h \circ f \circ g$ .

Then by Mori theorem,

$$|\phi(z_1) - \phi(z_2)| \leq 16|z_1 - z_2|^{\frac{1}{k}}.$$

And since  $E$  is compact.

Let  $|g^{-1}(z)| \leq r_0 < 1$  for  $z \in E$ .

Then by Koebe distortion theorem,

$$\begin{aligned} |g(z_1) - g(z_2)| &\geq \frac{|g'(z_1)|}{4} \left| \frac{z_1 - z_2}{1 - z_1 \bar{z}_2} \right| \\ &\geq \frac{(1 - |z_1|)|g'(0)|}{64} |z_1 - z_2| \\ &\geq \frac{(1 - r_0)(1 - |\alpha|)^2}{64|\alpha|} |p - q| |z_1 - z_2| \\ &\geq \frac{(1 - r_0)^3}{64|\alpha|} |p - q| |z_1 - z_2| \end{aligned}$$

On the other hand, for  $|z| \leq r_0$ , by Mori theorem, we obtain

$$1 - r_0 = 1 - |z| \leq 16(1 - |\phi(z)|)^{\frac{1}{k}}, |\alpha| \leq 16|\beta|^{\frac{1}{k}}.$$

So  $|\phi(z)| \leq \rho_0 < 1$ , where  $\rho_0$  only depends on  $r_0$ .

So similarly, we have

$$|z_1 - z_2| \leq \frac{4}{(1 - \rho_0)^6 |\beta|} |p - q| |h(z_1) - h(z_2)|.$$

Hence

$$\begin{aligned} |f(z_1) - f(z_2)| &\leq \frac{4}{(1 - \rho_0)^6 |\beta|} |p - q| |h \circ f(z_1) - h \circ f(z_2)| \\ &\leq \frac{2^{6+4k}}{(1 - \rho_0)^6 |\alpha|^k} |p - q| \left( \frac{64|\alpha|}{(1 - r_0)^3 |p - q|} |z_1 - z_2| \right)^{\frac{1}{k}} \\ &\leq C|z_1 - z_2|^{\frac{1}{k}} \end{aligned}$$

where  $C$  only depends on  $E$ .  $\square$

## 1.6 quasiconformal maps on Riemann surface

**Def 1.6.1.**  $S_1, S_2$  are Riemann surface, then  $f : S_1 \rightarrow S_2$  is  $k$ -quasiconformal if  $\tilde{f} : \mathbb{D} \rightarrow \mathbb{D}$  is  $k$ -quasiconformal.

**Lemma 1.6.1.** Suppose  $f : S_1 \rightarrow S_2$  is a  $k$ -quasiconformal map and  $\gamma_1 \subset S_1$  a closed geodesic on  $S_1$ , let  $\gamma_2 \subset S_2$  be the closed geodesic which is homotopic to  $f(\gamma_1)$ , then  $l(\gamma_2) \leq kl(\gamma_1)$ .

*Proof.* Consider  $f_* : \pi_1(S_1, *) \rightarrow \pi_1(S_2, *)$ .

After identifying  $\pi_1$  with  $\Gamma_i \subset \text{Aut}(\mathbb{D})$ , we obtain a lift  $\tilde{f} : \mathbb{D} \rightarrow \mathbb{D}$  such that

$$\tilde{f} \circ A_1 = A_2 \circ \tilde{f}, f_*(A_1) = A_2.$$

Let  $\gamma_i$  be the axis of  $A_i$ .

We first assume  $\tilde{f}(p) \in \gamma_2$  where  $p \in \gamma_1$ , then

$$\begin{aligned} nl(\gamma_2) &\leq d\left(\tilde{f}(p), A_2^n\left(\tilde{f}(p)\right)\right) = d\left(\tilde{f}(p), \tilde{f}(A_1^n(p))\right) \\ &\leq kd(p, A_1^n(p)) + C = knl(\gamma_1) + C. \end{aligned}$$

So  $l(\gamma_2) \leq kl(\gamma_1) + \frac{C}{n}$ , i.e.  $l(\gamma_2) \leq kl(\gamma_1)$  as  $n \rightarrow \infty$ .

For the general case, take a fixed point  $q \in \gamma_2$ , then

$$\left|d\left(\tilde{f}(p), A_2^n\left(\tilde{f}(p)\right)\right) - d(q, A_2^n(q))\right| \leq 2d(\tilde{f}(p), q) = C_1.$$

□

**Prop 1.6.1.** There exists no quasiconformal map  $f : \mathbb{D} \rightarrow \mathbb{C}$ .

*Proof.* Let  $\Omega = \{|z| \leq r_0 < 1\}$  and  $f(\Omega) \subset \{|z| \leq R\}$ .

Then by definition,

$$m(\mathbb{C} \setminus f(\Omega)) \leq km(\{r_0 < |z| < 1\}) = \frac{1}{2\pi} \log \frac{1}{r_0}.$$

On the other hand, for arbitrary  $M$ ,

$$m(\mathbb{C} \setminus B_R) \geq m(\{R \leq |z| \leq M\}) = \frac{1}{2\pi} \log \frac{M}{R}.$$

Take  $M \geq \frac{R}{r_0}$ , contradiction!

□

## 1.7 Topological definition of quasiconformal maps

**Def 1.7.1.**  $f : \mathbb{S}^2 \rightarrow \mathbb{S}^2$  is a homeomorphism, define

$$N_f = \{A \circ f \circ B \mid A, B \in \text{Aut}(\mathbb{S}^2), A \circ f \circ B \text{ fixes } p, q, r \in \mathbb{S}^2\},$$

$$\text{Homeo}(\mathbb{S}^2, p, q, r) = \{g : \mathbb{S}^2 \rightarrow \mathbb{S}^2 \mid g \text{ is homeomorphism, } g \text{ fixes } p, q, r\}.$$

**Lemma 1.7.1.** If  $f$  is quasiconformal, then  $N_f$  is sequentially precompact in  $\text{Homeo}(\mathbb{S}^2, p, q, r)$ .

*Proof.* Consider the stereographic projection  $g : \mathbb{S}^2 \rightarrow \bar{\mathbb{C}}$  with  $g(p) = \infty$ .

Then for a sequence  $\{A_n \circ f \circ B_n\} \subset N_f$ ,  $g \circ A \circ f \circ B \circ g^{-1}$  is a  $k$ -quasiconformal map on  $\mathbb{C}$ .

So  $\{g \circ A_n \circ f \circ B_n \circ g^{-1}\}$  converges to a  $k$ -quasiconformal map  $f_0 : \mathbb{C} \rightarrow \mathbb{C}$ .

Take  $h = g^{-1} \circ f_0 \circ g$ ,  $h(p) = p$ .

Hence  $h$  is a homeomorphism in  $\text{Homeo}(\mathbb{S}^2, p, q, r)$ .

□

**Exam 1.7.1.** Take

$$f(z) = \left(e^{|z|} - 1\right) e^{i \arg(z)}, B_n(z) = nz, A_n(z) = \frac{z}{e^n - 1}.$$

Then  $A_n \circ f \circ B_n$  fixes  $0, 1, \infty$  but maps  $2$  to

$$\frac{e^{2n} - 1}{e^n - 1} \rightarrow \infty$$

**Lemma 1.7.2.** If  $f$  is not quasiconformal, then for any  $n > 0$ , there exists quadrilateral  $Q_n$  with  $m(Q_n) = 1$  such that  $nm(f(Q_n)) < 1$

*Proof.* Since  $f$  is not quasiconformal.

Take quadrilateral  $Q$  such that  $2nm(f(Q)) < m(Q)$ .

WLOG, we assume  $Q$  is a rectangle and  $m(Q) \geq 1$ , otherwise we swap the a-sides with b-sides.

If  $m(Q) = 1$ , then there is nothing need to be proved, so we only consider  $m(Q) > 1$ .

Let  $k < m(Q) \leq k+1$  where  $k \in \mathbb{Z}^+$  and divide  $Q$  into  $k$  squares  $R_1, \dots, R_k$  and a rectangle  $R_{k+1}$ , which may also be a square, then

$$m(Q) > 2nm(f(Q)) \geq 2n \sum_{i=1}^{k+1} m(f(R_i)) > 2n \sum_{i=1}^k m(f(R_i)).$$

So there exists some  $i \leq k$  such that

$$m(f(R_i)) < \frac{m(Q)}{2nk} \leq \frac{k+1}{2nk} \leq \frac{1}{n}.$$

Hence take  $Q_n = R_i$ , we have  $nm(f(Q_n)) < 1$  and  $m(Q) = m(R_i) = 1$ . □

**Thm 1.7.1** (Topological definition). A homeomorphism  $f : \mathbb{S}^2 \rightarrow \mathbb{S}^2$  is quasiconformal iff  $N_f$  is sequentially precompact subset of  $\text{Homeo}(\mathbb{S}^2, p, q, r)$ .

*Proof.* The “only if” part is proved in lemma 1.7.1, we now prove the “if” part.

WLOG, we assume  $p, q, r$  are  $0, 1, \infty$  resp.

For  $z \in \mathbb{C}$ , take  $x, y \in \partial B_r(z)$  and  $\hat{f} = A \circ f \circ B \in N_f$  such that

$$B(0) = z, B(1) = x, A(f(z)) = 0, A(f(x)) = 1, B(\infty) = A(\infty) = \infty.$$

Since  $A, B$  are affine map, we have

$$d(0, B^{-1}(y)) = 1, \frac{d(f(y), f(z))}{d(f(x), f(z))} = d(0, A(f(y))).$$

Suppose  $f$  is not quasiconformal, then there exists  $z_n, x_n, y_n$  such that

$$|B_n^{-1}(y_n)| = 1, \left| \hat{f}_n(B_n^{-1}(y_n)) \right| > n.$$

WLOG, assume  $\hat{f}_n$  converges to  $g \in \text{Homeo}(\mathbb{S}^2, 0, 1, \infty)$  and  $B_n^{-1}(y_n)$  converges to  $y$ .

Then  $|g(y)| \geq \lim_{n \rightarrow \infty} n = \infty$ .

But  $g(\infty) = \infty$  and  $g$  is injective, contradiction! □

## 1.8 Analytical properties of quasiconformal maps

**Def 1.8.1.** Given a quasiconformal map  $f : U \rightarrow f(U)$ , we say  $f$  is absolutely continuous on lines if in every rectangle  $R \subset U$ ,  $f$  is absolutely continuous on almost every horizontal and almost every vertical line.

**Lemma 1.8.1.** *Let  $f : U \rightarrow f(U)$  be quasiconformal, then  $f$  is absolutely continuous on lines.*

*Proof.* Take disjoint intervals  $(a_1, b_1), \dots, (a_n, b_n)$  in  $[\alpha, \beta]$ .

Let  $A(y)$  be the image area under  $f$  of the rectangle  $[\alpha, \beta] \times [0, y]$ .

Then  $A$  is increasing, *i.e.* derivative  $A'(y)$  exists a.e.

WLOG, assume  $A'(y_0)$  exists.

Consider  $Q = [\alpha, \beta] \times [y_0, y_0 + \delta]$ ,  $Q_i = [a_i, b_i] \times [y_0, y_0 + \delta]$  and  $Q' = f(Q)$ ,  $Q'_i = f(Q_i)$ .

Let  $\gamma_0$  be the line connecting  $a_i$  and  $b_i$ ,  $l_i = l(\gamma_0) = |b_i - a_i|$ ,  $l'_i = l(f(\gamma_0))$ .

We first show that for sufficiently small  $\delta$ , the length of any arc  $\gamma$  connecting the b-side of  $Q'_i$  is near by  $l'_i$ .

Take a partition  $a_i = t_0 < t_1 < \dots < t_m = b_i$  and  $\zeta_i = f(t_k, y_0)$  such that

$$\sum_{k=1}^m |\zeta_k - \zeta_{k-1}| \geq l'_i - \frac{\varepsilon}{2}.$$

Take sufficiently small  $\delta$  such that the variation of  $f$  on vertical segments  $\{t_k\} \times [y_0, y_0 + \delta]$  is less than  $\frac{\varepsilon}{4n}$ , then we have

$$l(\gamma) \geq \sum_{k=1}^m |\zeta_k - \zeta_{k-1}| - \frac{\varepsilon}{2} \geq l'_i - \varepsilon$$

So take  $\varepsilon < \frac{1}{2} \min\{l'_i\}$ , we obtain

$$\frac{(l'_i)^2}{4A(Q_i)} \leq m_i(Q'_i) \leq k \frac{l_i}{\delta},$$

$$\left( \sum_{i=1}^n l'_i \right)^2 \leq \sum_{i=1}^n \frac{(l'_i)^2}{l_i} \cdot \sum_{i=1}^n l_i \leq 4K \frac{A(y_0 + \delta) - A(y_0)}{\delta} \sum_{i=1}^n l_i.$$

And since

$$\lim_{\delta \rightarrow 0} \frac{A(y_0 + \delta) - A(y_0)}{\delta} = A'(y_0) < +\infty,$$

Hence as  $\sum l_i \rightarrow 0$ ,  $\sum l'_i \rightarrow 0$ , *i.e.*  $f$  is absolutely continuous on horizontal line.

Similarly,  $f$  is absolutely continuous on vertical line. □

**Thm 1.8.1.** *Let  $f : U \rightarrow f(U)$  be quasiconformal, then  $f$  is differentiable a.e. on  $U$  and it is differentiable in the sense of distribution.*

*Proof.* By lemma 1.8.1,  $f$  is absolutely continuous on lines.

So  $f$  has partial derivatives  $f_x, f_y$  a.e.

By Egoroff theorem, the limits

$$f_x(z) = \lim_{\delta \rightarrow 0} \frac{f(z + \delta) - f(z)}{\delta}, f_y(z) = \lim_{\delta \rightarrow 0} \frac{f(z + \delta i) - f(z)}{\delta}.$$

are taken uniformly except on  $U \setminus E$  of arbitrary small measure.

Then it is sufficient to prove that  $f$  is differentiable a.e. on  $E$ .

Notice that almost every point  $x_0 + y_0 i \in E$  is point of density for  $E \cap (\{x = x_0\} \cup \{y = y_0\})$ .

So it is sufficient to prove that  $f$  is differentiable at such a point  $z_0 = x_0 + y_0i$ .  
And WLOG, we assume  $z_0 = 0$ .

Then there exists  $\delta > 0$  such that for  $|x|, |y|, |\delta_1|, |\delta_2| < \delta$ ,

$$|f_x(z) - f_x(0)| < \varepsilon, |f_y(z) - f_y(0)| < \varepsilon,$$

$$\left| \frac{f(z + \delta_1) - f(z)}{\delta_1} - f_x(z) \right| < \varepsilon, \left| \frac{f(z + \delta_2 i) - f(z)}{\delta_2} - f_y(z) \right| < \varepsilon.$$

So

$$f(z) - f(0) - x f_x(0) - y f_y(0) = (f(z) - f(x) - y f_y(x)) + (f(x) - f(0) - x f_x(0)) + y (f_y(x) - f_y(0))$$

If  $x \in E$  or  $y \in E$ , we have

$$|f(z) - f(0) - x f_x(0) - y f_y(0)| \leq 3\varepsilon|z|.$$

Now we want to prove the case when  $x, y \notin E$ .

Since  $\frac{m(E \cap (-x, x))}{2|x|} \rightarrow 1$  as  $x \rightarrow 0$ .

Take  $\delta$  sufficiently small such that for  $x < \delta$

$$m(E \cap (-x, x)) > \frac{2 + \varepsilon}{1 + \varepsilon}|x|.$$

Then  $E \cap \left(\frac{x}{1 + \varepsilon}, x\right) \neq \emptyset$ .

Similarly, we apply this process to  $y$ .

So if  $|z| < \frac{\delta}{1 + \varepsilon}$ , there exists  $x_1, x_2, y_1, y_2 \in E$  such that

$$\frac{x}{1 + \varepsilon} < x_1 < x < x_2 < (1 + \varepsilon)x, \frac{y}{1 + \varepsilon} < y_1 < y < y_2 < (1 + \varepsilon)y.$$

Consider rectangle  $(x_1, x_2) \times (y_1, y_2)$ .

By the maximal principle, there exists a point  $z^*$  on the perimeter such that

$$\begin{aligned} |f(z) - f(0) - x f_x(0) - y f_y(0)| &\leq |f(x^* + iy^*) - f(0) - x f_x(0) - y f_y(0)| \\ &\leq 3\varepsilon|z^*| + |x - x^*||f_x(0)| + |y - y^*||f_y(0)| \\ &\leq 3\varepsilon(1 + \varepsilon)|z| + \varepsilon|f_x(0)||z| + \varepsilon|f_y(0)||z| \end{aligned}$$

Hence  $f$  is differentiable a.e.

Moreover, for compact  $K \subset U$ , we have

$$\int_K J_f dx dy \leq A(f(K)).$$

And since  $f$  is  $k$ -quasiconformal.

Similar to proposition 1.4.2, we can prove that

$$|f_{\bar{z}}| \leq K|f_z|, |f_z|^2 \leq \frac{J}{1 - K^2} \text{ with } K = \frac{k - 1}{k + 1}.$$

So  $f_x, f_y$  are locally square integrable.

Take a test function  $\phi$ , which is  $C^1$  with compact support.

By the integration over horizontal lines and Fubini theorem,

$$\iint f_x \phi dx dy = - \iint f \phi_x dx dy, \iint f_y \phi dx dy = - \iint f \phi_y dx dy.$$

□

## Chapter 2

# Boundary Correspondence

### 2.1 Quasi-isometry maps

**Def 2.1.1.**  $E$  is a convex subset of  $\mathbb{H}^n$ , then one can define the nearest point retraction map  $\pi : \mathbb{H}^n \rightarrow E$ : for  $p \in \mathbb{H}^n$ , there exists a unique point  $q \in E$  such that  $d(p, E) = d(p, q)$  and we set  $\pi(p) = q$ .

**Lemma 2.1.1.**  $\|d\pi(z)\| \leq \cosh(r)^{-1}$ , where  $r = d(z, E)$ .

*Proof.* Let  $r = d(z, E)$  and consider a smooth curve  $\gamma$  in the level set

$$\Sigma_r = \{w \in \mathbb{H}^n | d(w, E) = r\}.$$

with  $\gamma(0) = z, \gamma'(0) = v \in T_z \Sigma_r$ .

Then  $\pi \circ \gamma$  is a curve in  $E$  with  $(\pi \circ \gamma)'(0) = d\pi_z(v)$ .

Let  $\alpha(t, -)$  be the unit-speed geodesic from  $\pi(\gamma(t))$  to  $\gamma(t)$  and Consider the Jacobi field

$$J(s) = \alpha \left( \frac{\partial}{\partial t} \right) \Big|_{t=0}.$$

Then  $\alpha(t, -)$  is perpendicular to  $\partial E$  and  $J(0) = d\pi_z(v), J(r) = v$ .

And since  $\mathbb{H}^n$  has constant curvature  $-1$ , so we obtain

$$J(s) = d\pi(z)(v) \cosh(s) + J'(0) \sinh(s).$$

On the other hand,

$$J'(0) = \hat{\nabla}_{\frac{\partial}{\partial s}} \alpha_* \left( \frac{\partial}{\partial t} \right) \Big|_{s=t=0} = \hat{\nabla}_{\frac{\partial}{\partial t}} \alpha_* \left( \frac{\partial}{\partial s} \right) \Big|_{s=t=0} = \hat{\nabla}_{\frac{\partial}{\partial t}} N \Big|_{t=0} = \nabla_{J(0)} N$$

where  $N$  is the unit normal vector to  $E$  pointing outward.

Notice that  $E$  is convex.

So the second fundamental form

$$B(J(0), J(0)) = \langle \nabla_{J(0)} N, J(0) \rangle = \langle J'(0), J(0) \rangle \geq 0.$$

Therefore we have

$$\begin{aligned} |v|^2 &= |J(r)|^2 \\ &= \cosh^2(r) |d\pi_z(v)|^2 + \sinh^2(r) |J'(0)|^2 + 2 \cosh(r) \sinh(r) \langle d\pi_z(v), J'(0) \rangle \\ &\geq \cosh^2(r) |d\pi_z(v)|^2 \end{aligned}$$

And since  $d\pi_z$  vanishes in the direction perpendicular to  $\Sigma_r$ .

Hence  $\|d\pi(z)\| \leq \cosh(r)^{-1}$ . □

**Def 2.1.2.** A map  $\gamma : \mathbb{R} \rightarrow \mathbb{H}^n$  is called  $(L, A)$ -quasigeodesic if

$$\frac{|x - y|}{L} - A \leq d(\gamma(x), \gamma(y)) \leq L|x - y| + A.$$

**Lemma 2.1.2.** For  $L, A > 0$ , there exists  $A_1, L_1, D_1$  depending on  $(L, A)$  such that for every  $(L, A)$ -quasigeodesic  $\gamma$ , there exists a  $(L, A_1)$ -quasigeodesic  $\gamma_1$  such that

(1)  $l_{\mathbb{H}^n}(\gamma_1([a, b])) \leq L_1|a - b|.$

(2)  $d_{\mathbb{H}^n}(\gamma(x), \gamma_1(x)) \leq D_1$  for every  $x$ .

*Proof.* Let  $\gamma_1|_{[n, n+1]}$  be the geodesic from  $\gamma(n)$  to  $\gamma(n+1)$  with  $|\gamma'(t)| \equiv C$ .

Then for  $n \leq a < n+1, m \leq b < m+1$ ,

$$\begin{aligned} & l_{\mathbb{H}^n}(\gamma_1([a, b])) \\ &= l_{\mathbb{H}^n}(\gamma_1([a, n+1])) + \sum_{i=n+1}^{m-1} l_{\mathbb{H}^n}(\gamma_1([i, i+1])) + l_{\mathbb{H}^n}(\gamma_1[m, b]) \\ &= (n+1-a)d(\gamma(n), \gamma(n+1)) + \sum_{i=n+1}^{m-1} d(\gamma(i), \gamma(i+1)) + (b-m)d(\gamma(m), \gamma(m+1)) \\ &\leq (L+A)|b-a| \end{aligned}$$


And for  $n \leq x < n+1$ ,

$$d(\gamma(x), \gamma_1(x)) \leq d(\gamma(x), \gamma(n)) + d(\gamma_1(n), \gamma_1(x)) \leq 2(L+A).$$

□

**Thm 2.1.1** (Morse lemma). Suppose  $\gamma : I \rightarrow \mathbb{H}^n$  is an  $(L, A)$ -quasigeodesic where  $I$  is an interval either finite or infinite, then  $\gamma(I)$  is within a bounded distance  $R$  from a geodesic in  $\mathbb{H}^n$ , where  $R$  only depends on  $L$  and  $A$ .

*Proof.*



by lemma 2.1.2 we can assume  $\gamma$  is  $(L, A)$ -quasigeodesic and  $l(\gamma([a, b])) \leq L|a - b|.$

We first consider the case that  $I = [T_1, T_2]$  is finite.

Let  $\delta$  be the geodesic arc between  $\gamma(T_1)$  and  $\gamma(T_2)$  and  $(a, b) \subset [T_1, T_2]$  be a maximal interval such that  $\gamma((a, b))$  lies outside the cylinder  $B(\delta, r_0)$  where

$$\frac{2r_0L^2 + LA}{\cosh(r_0) - L^2} \leq 1.$$

Then we have

$$\begin{aligned} \cosh(r_0)d(\pi(\gamma(a)), \pi(\gamma(b))) &\leq l(\gamma(a, b)) \leq L|a - b| \\ &\leq L(Ld(\gamma(a), \gamma(b)) + A) \\ &\leq L^2(2r_0 + d(\pi(\gamma(a)), \pi(\gamma(b)))) + LA \end{aligned}$$

So

$$d(\pi(\gamma(a)), \pi(\gamma(b))) \leq \frac{2r_0L^2 + LA}{\cosh(r_0) - L^2} \leq 1.$$

Therefore  $d(\gamma(a), \gamma(b)) \leq \cosh(r_0)$ , i.e.  $|a - b| \leq L \cosh(r_0) + A$ .

Replace  $r_0$  by  $r_0 + L(L \cosh(r_0) + A) + A$ , then  $\gamma([T_1, T_2]) \subset B(\delta, r_0)$ .

For general case, take a sequence of finite interval  $I_1 \subset I_2 \subset \dots \subset I$  and let  $\delta_i$  be the corresponding geodesic arc.

Notice that  $\delta_i$  are all  $L$ -Lipschitz.

By Ascoli-Arzela theorem and diagonal argument,  $\delta_i$  converges to a geodesic  $\delta$  as  $i \rightarrow \infty$

Hence as  $i \rightarrow \infty$ ,  $\gamma(I) \subset B(\delta, r_0)$ .  $\square$

**Def 2.1.3.** A map  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$  is called  $(L, A)$ -quasi-isometry if

$$\frac{d(x, y)}{L} - A \leq d(f(x), f(y)) \leq Ld(x, y) + A,$$

**Thm 2.1.2.**  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$  is a quasi-isometry then  $f$  extend continuous to a quasiconformal map  $\partial f : \partial\mathbb{H}^n \rightarrow \partial\mathbb{H}^n$ .

We can define  $\partial f$  directly by Morse lemma, and to prove this theorem, we need some deeper properties of quasi-isometry maps. For brevity, we have placed the proof in chapter A.

## 2.2 $M$ -condition

**Def 2.2.1.** Let  $h : \mathbb{R} \rightarrow \mathbb{R}$  be a homeomorphism, we say that  $h$  satisfies the  $M$ -condition if

$$\frac{1}{M} \leq \frac{h(x+t) - h(x)}{h(x) - h(x-t)} \leq M.$$

Such  $h$  is also called quasisymmetric.

**Lemma 2.2.1.** Let  $f : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  be  $k$ -quasiconformal such that  $f(\infty) = \infty$ , then  $h(x) = f(x, 0)$  is an increasing homeomorphism which satisfies the  $M$ -condition for same  $M = M(k)$ .

*Proof.* For  $t \in \mathbb{R}$ , By choosing affine map  $A, B$ , let  $g = B \circ f \circ A$  such that

$$g(-1) = -1, g(0) = 0, g(1) = \frac{h(x+t) - h(x)}{h(x) - h(x-t)}.$$

Suppose there exists  $f_m : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  such that

$$\frac{h_m(x_m + t_m) - h_m(x_m)}{h_m(x_m) - h_m(x_m - t_m)} \rightarrow 0 \text{ or } \infty$$

Then we can rescaled  $f_n$  to get  $g_n$  such that

$$g_n(-1) = -1, g_n(0) = 0, g_n(1) \rightarrow 0 \text{ or } \infty.$$

And since  $g_n$  fixed  $0, -1, \infty$ .

So  $g_n$  converge to a  $k$ -quasiconformal  $g_n \rightarrow g$ .

But  $g(1) = \lim_{n \rightarrow \infty} g_n(1) = 0 \text{ or } \infty$ , contradiction!  $\square$

**Lemma 2.2.2.** Let  $h : \mathbb{R} \rightarrow \mathbb{R}$  be a homeomorphism satisfying  $M$ -condition with  $h(0) = 0$ ,  $h(1) = 1$ , then

$$\frac{1}{M+1} \leq \int_0^1 h(x) dx \leq \frac{M}{M+1}.$$

*Proof.* Let  $F(x) = \sup h_0(x)$  for all quasi-isometry  $h_0$  with  $h_0(0) = 0, h_0(1) = 1$ .

Then for  $0 < t < 1$ ,

$$\frac{h\left(\frac{t}{2}\right)}{h(t)} \leq F\left(\frac{1}{2}\right)$$

So

$$F\left(\frac{t}{2}\right) \leq F\left(\frac{1}{2}\right)F(t).$$

And similarly, we have

$$\frac{h\left(\frac{1-t}{2} + t\right) - h(t)}{1 - h(t)} \leq F\left(\frac{1}{2}\right).$$

Therefore

$$F\left(\frac{1+t}{2}\right) \leq F\left(\frac{1}{2}\right) + \left(1 - F\left(\frac{1}{2}\right)\right)F(t).$$

Thus we obtain

$$F\left(\frac{t}{2}\right) + F\left(\frac{1+t}{2}\right) \leq F\left(\frac{1}{2}\right) + F(t).$$

Hence

$$\begin{aligned} \int_0^1 h(t)dt &\leq \int_0^1 F(t)dt = \frac{1}{2} \int_0^2 F\left(\frac{t}{2}\right) dt \\ &= \frac{1}{2} \int_0^1 \left(F\left(\frac{t}{2}\right) + F\left(\frac{1+t}{2}\right)\right) dt \\ &\leq \frac{1}{2}F\left(\frac{1}{2}\right) + \frac{1}{2} \int_0^1 F(t)dt \\ &\leq F\left(\frac{1}{2}\right) \leq \frac{M}{M+1} \end{aligned}$$

□

**Lemma 2.2.3** (Gauge rule). *Let  $f : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  be  $k$ -quasiconformal with  $f(\infty) = \infty$ , then for any affine map  $A, B$ , take  $h(x) = f(x, 0), h_1(x) = (B \circ f \circ A)(x, 0)$ , prove that*

$$h_1 = B \circ h \circ A.$$

*Proof.* Let  $A = az + b, B = cz + d$  with  $a, c > 0$  and  $b, d \in \mathbb{R}$ , then

$$(B \circ f \circ A)(x, 0) = cf(ax + b, 0) + d = ch(ax + b) + d = B \circ h \circ A.$$

□

**Thm 2.2.1.** *Let  $h : \mathbb{R} \rightarrow \mathbb{R}$  be a homeomorphism satisfying  $M$ -condition, then there exists a map  $\phi = u + iv$  which is  $k$ -quasiconformal for  $k = 2M(M + 1)$  and  $\phi : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  extending  $h$ .*

*Proof.* Take

$$u(x, y) = \frac{1}{2y} \int_{-y}^y h(x+t)dt, v(x, y) = \frac{1}{2y} \int_0^y (h(x+t) - h(x-t))dt.$$

Then  $v(x, y) \geq 0$  and  $v(x, y) \rightarrow 0$  as  $y \rightarrow 0$ , i.e.  $\phi$  is well-defined and  $\phi(x, 0) = u(x, 0) = h(x)$ .

And since we have

$$u_x = \frac{1}{2y}(h(x+y) - h(x-y)),$$

$$\begin{aligned}
 u_y &= -\frac{1}{2y} \int_{x-y}^{x+y} h dt + \frac{1}{2y} (h(x+y) + h(x-y)), \\
 v_x &= \frac{1}{2y} (h(x+y) - 2h(x) + h(x-y)), \\
 v_y &= -\frac{1}{2y^2} \left( \int_x^{x+y} h dt - \int_{x-y}^x h dt \right) + \frac{1}{2y} (h(x+y) - h(x-y)).
 \end{aligned}$$

By gauge rule, we can assume  $h(0) = 0, h(1) = 1$  and we only need to compute dilatation at  $z = i$ , then

$$\begin{aligned}
 u_x &= \frac{1}{2} (1 - h(-1)), u_y = -\frac{1}{2} \int_{-1}^1 h dt + \frac{1 + h(-1)}{2}, \\
 v_x &= \frac{1 + h(-1)}{2}, v_y = -\frac{1}{2} \left( \int_0^1 h dt - \int_{-1}^0 h dt \right) + \frac{1}{2} (1 - h(-1)).
 \end{aligned}$$

Let

$$\xi = 1 - \int_0^1 h dt, \beta = -h(-1), \eta\beta = -h(-1) + \int_{-1}^0 h dt.$$

So we obtain

$$u_x = \frac{1 + \beta}{2}, v_x = \frac{1 - \beta}{2}, u_y = \frac{\xi - \eta\beta}{2}, v_y = \frac{1}{2} (\xi + \eta\beta).$$

$$\begin{aligned}
 d^2 &= \left| \frac{((1 - \xi) + \beta(1 - \eta)) + i((1 + \xi) - \beta(1 + \eta))}{((1 + \xi) + \beta(1 + \eta)) + i((1 - \xi) - \beta(1 - \eta))} \right|^2 \\
 &= \frac{1 + \xi^2 + \beta^2(1 + \eta^2) - 2\beta(\xi + \eta)}{1 + \xi^2 + \beta^2(1 + \eta^2) + 2\beta(\xi + \eta)}
 \end{aligned}$$

Since we have estimates

$$M^{-1} \leq \beta \leq M, \frac{1}{M+1} \leq \xi, \eta \leq \frac{M}{M+1}.$$

Therefore

$$D = \frac{1 + d}{1 - d} \leq 2 \frac{1 + d^2}{1 - d^2} < 2M(M+1).$$

Moreover, we must show that  $\phi(z) \rightarrow \infty$  as  $z \rightarrow \infty$ , this is because

$$u^2 + v^2 = \frac{1}{2y} \left( \left( \int_x^{x+y} h dt \right)^2 + \left( \int_{x-y}^x h dt \right)^2 \right).$$

Hence by monodromy theorem,  $\phi$  is homeomorphic, *i.e.*  $\phi$  is  $k$ -quasiconformal.  $\square$

**Prop 2.2.1.** *X and Y are Riemann surface diffeomorphic to  $\Sigma_g$ , then they are biholomorphic iff they are isometric w.r.t. hyperbolic metrics.*

**Prop 2.2.2.** *X and Y are Riemann surface diffeomorphic to  $\Sigma_g$  but not biholomorphic, let  $f : X \rightarrow Y$  be a diffeomorphism and  $g : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  is the lift of  $f$ , then  $g$  is not differentiable anywhere with non-zero derivative on the boundary.*

*Proof.* Let  $\Gamma_X, \Gamma_Y$  be the Fuchian groups.

Then for  $A \in \Gamma_X$ , there exists  $B \in \Gamma_Y$  such that  $g \circ A = B \circ g$ .

WLOG, assume  $g(0) = 0$  and suppose  $g$  is differentiable at 0 with non-zero derivative.

Let  $A_n(z) = \frac{z}{n}, g_n = A_n^{-1} \circ g \circ A_n$ .

Since  $g'(0) \neq 0$ .

Take arbitrary  $z_0 \in \mathbb{H}^2$  and a fundamental domain  $F$  of  $X$ .

Then there exists  $\gamma_n \in \Gamma_X$  such that  $\gamma_n \circ A_n(z_0) \in F$ .

So by Ascoli-Arzela theorem, there exists  $\{n_i\}$  such that  $\gamma_{n_i} \circ A_{n_i}$  converges to  $\sigma_1 : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ .

Therefore  $A_{n_i}^{-1} \circ \gamma_{n_i}^{-1} \circ \gamma \circ \gamma_{n_i} \circ A_{n_i}$  converges to some  $\sigma_1^{-1} \gamma \sigma_1$  for every  $\gamma \in \Gamma_X$ .

And consider  $\delta_n \in \Gamma_Y$  such that  $g \circ \gamma_n = \delta_n \circ g$ .

Then  $\delta_n A_n \left( ng \left( \frac{z_0}{n} \right) \right) \in g(F)$ .

WLOG, we assume  $\delta_{n_i} \circ A_{n_i}$  converges to  $\sigma_2 : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ .

Let  $g_n$  converges to  $h : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  and  $a = g'(0)$ , then

$$\begin{aligned} h \circ \sigma_1^{-1} \circ \gamma \circ \sigma_1 &= \lim_{i \rightarrow \infty} g_{n_i} \circ A_{n_i}^{-1} \circ \gamma_{n_i}^{-1} \circ \gamma \circ \gamma_{n_i} \circ A_{n_i} \\ &= \lim_{i \rightarrow \infty} A_{n_i}^{-1} \circ g \circ \gamma_{n_i}^{-1} \circ \gamma \circ \gamma_{n_i} \circ A_{n_i} \\ &= \lim_{i \rightarrow \infty} A_{n_i}^{-1} \circ \delta_{n_i}^{-1} \circ \delta \circ \delta_{n_i} \circ g \circ A_{n_i} \\ &= \sigma_2^{-1} \circ \delta \circ \sigma_2 \circ h \end{aligned}$$

where  $g \circ \gamma = \delta \circ g$  for  $\gamma \in \Gamma_X$ ,  $\delta \in \Gamma_Y$ .

Notice that for  $x \in \mathbb{R}$ ,

$$h(x) = \lim_{n \rightarrow \infty} \frac{g\left(\frac{x}{n}\right) - g(0)}{\frac{1}{n}} = ax$$

Hence we can get  $h : X \rightarrow Y, x \mapsto ax$  which is linear and biholomorphic, contradiction!  $\square$

**Coro 2.2.1.** *There are quasisymmetric map  $h : \mathbb{R} \rightarrow \mathbb{R}$  which are not absolutely continuous.*

*Proof.* Consider the biholomorphic map  $g : \mathbb{H}^2 \rightarrow \mathbb{H}^2$  in proposition 2.2.2.

Then  $\partial g$  is not absolutely continuous.  $\square$

Using similar method, we can prove the Mostow rigidity theorem. Since the proof is so long and is not quite related to the course, it can be found in chapter A.

**Thm 2.2.2** (Mostow rigidity).  *$M, N$  are closed hyperbolic  $n$ -manifold with  $n \geq 3$ , then every homotopy equivalent  $f : M \rightarrow N$  must be homeomorphism to an isometry.*

**Lemma 2.2.4.** *Let  $X$  be a closed Riemann surface with genus  $g \geq 2$ , then there is a unique closed geodesic in a free homotopy type w.r.t. parametrization.*

*Proof.* Consider the covering map  $\pi : \mathbb{H}^2 \rightarrow X$ , a free homotopy type in  $X$  corresponds to a deck transformation  $F : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ .

For a closed geodesic  $\gamma$  in the given free homotopy type with  $\gamma(1) = \gamma(0)$ .

Consider the lifting  $\tilde{\gamma} : \mathbb{R} \rightarrow \mathbb{H}^2$ .

Then  $\tilde{\gamma}(t+1) = F(\tilde{\gamma}(t))$ , i.e.  $\tilde{\gamma}$  is the axis of  $F$ .

Hence  $\gamma$  is unique w.r.t. parametrization.  $\square$

**Lemma 2.2.5.** *Let  $X$  be a closed Riemann surface with genus  $g \geq 2$  and  $f$  be an automorphism that homotopies to Id, then  $f = \text{Id}$ .*

*Proof.* Consider a figure-eight closed geodesic  $\gamma$  that self-intersects at  $p$ .

Since  $f \simeq \text{Id}$ .

So  $f(\gamma)$  and  $\gamma$  are in the same free homotopy type and are both closed geodesic.

By lemma 2.2.4,  $f(\gamma) = \gamma$ , in particular,  $f(p) = p$ .

Denote the two parts of  $\gamma$  divided by  $p$  by  $\gamma_1, \gamma_2$ , which are geodesics.

We only need to prove that  $f(\gamma_i) = \gamma_i$  since  $f$  is an isometry.

Suppose  $f(\gamma_1) = \gamma_2, f(\gamma_2) = \gamma_1$ .

Then  $[\gamma_2] \cdot [\gamma_1] = [f(\gamma)] = [\gamma] = [\gamma_1] \cdot [\gamma_2]$ .

By Preissmann theorem,  $[\gamma_1] = [a]^p$  and  $[\gamma_2] = [a]^q$  for some  $[a] \in \pi_1(X)$  and  $p, q \in \mathbb{Z}$ .

So  $[\gamma] = [a]^{p+q}$ , in this case,  $\gamma$  must be  $(p+q)$ -times iteration of  $[a]$ , contradiction!  $\square$

**Prop 2.2.3.** Fix an closed Riemann surface  $X$  and let  $G$  be the group of all conformal automorphism of  $X$ , then  $|G| < +\infty$ .

*Proof.* Suppose  $|G| = \infty$ , let  $\{f_n\}$  be a sequence of automorphism.

Then  $f_n \rightarrow f \in G$ .

Let  $g_n = f_n^{-1} \circ f$ .

Then  $g_n \rightarrow \text{Id}$ .

So for sufficiently large  $n$ ,  $g_n$  is homotopic to  $\text{Id}$ , contradiction with lemma 2.2.5! □

**Def 2.2.2.** The restriction  $f : \mathbb{S}^1 \rightarrow f(\mathbb{S}^1)$  is called quasisymmetric map and  $f(\mathbb{S}^1)$  is called quasicircle.

# Chapter 3

## Beltrami differential

### 3.1 Beltrami differential

**Def 3.1.1.**  $f : \Omega \rightarrow f(\Omega)$  is a diffeomorphism, its Beltrami differential is defined as

$$\mu_f(z) = \text{Belt}(f)(z) = \left( \frac{f_{\bar{z}}}{f_z} \right) (z).$$

**Prop 3.1.1.**

$$D_f(z) = \frac{1 + |\mu_f(z)|}{1 - |\mu_f(z)|}.$$

*Proof.* Directly follows from proposition 1.1.2. □

**Prop 3.1.2.**

$$\mu_{h \circ f^{-1}} \circ f = \frac{f_z}{f_{\bar{z}}} \frac{\mu_h - \mu_f}{1 - \bar{\mu}_f \mu_h}.$$

*Proof.*

$$\begin{aligned} \frac{(h \circ f^{-1})_{\bar{z}}}{(h \circ f^{-1})_z}(f(z)) &= \frac{h_z(z) f_{\bar{z}}^{-1}(f(z)) + h_{\bar{z}}(z) \overline{f_z^{-1}(f(z))}}{h_z(z) f_z^{-1}(f(z)) + h_{\bar{z}}(z) \overline{f_{\bar{z}}^{-1}(f(z))}} \\ &= \frac{-h_z(z) \mu_f(z) + h_{\bar{z}}(z)}{h_z(z) - h_{\bar{z}}(z) \bar{\mu}_f} \cdot \frac{f_z - f_{\bar{z}} \bar{\mu}_f}{f_z(z) - \overline{f_{\bar{z}}(z)} \mu_f(z)} \\ &= \frac{\mu_h(z) - \mu_f(z)}{1 - \bar{\mu}_f \mu_h} \frac{f_z}{f_{\bar{z}}} \end{aligned}$$

□

**Coro 3.1.1.** If  $\mu_f = \mu_h$  a.e., then  $h \circ f^{-1}$  is conformal.

*Proof.*  $\mu_{h \circ f^{-1}} \circ f = 0$  a.e.

So  $h \circ f^{-1}$  is conformal. □

**Def 3.1.2.** A Riemannian metric  $ds^2$  on a differentiable surface is given by

$$ds^2 = E dx^2 + 2F dx dy + G dy^2$$

in  $(x, y)$  local coordinate. In other words,

$$ds = \lambda |dz + \mu d\bar{z}|$$

with

$$\lambda^2 = \frac{1}{4} \left( E + G + 2\sqrt{EG - F^2} \right), \mu = \frac{E - G + 2iF}{E + G + 2\sqrt{EG - F^2}}$$

Remark 3.1.1.

$$|\mu|^2 = \frac{E + G - 2\sqrt{EG - F^2}}{E + G + 2\sqrt{EG - F^2}} < 1.$$

**Prop 3.1.3.** Let  $f : \Omega \rightarrow \Omega_1$  such that  $\mu = \mu_f$ , then there exists  $\sigma(w)|dw|$  on  $\Omega_1$ , where  $w = f(z)$ , such that

$$f^*(\sigma(w)|dw|) = ds.$$

*Proof.*

$$\begin{aligned} f^*\sigma &= \sigma(f(z))|df| \\ &= (\sigma \circ f)(z)|f_z dz + f_{\bar{z}} d\bar{z}| \\ &= (\sigma \circ f)(z)|f_z|(z) \cdot |dz + \mu d\bar{z}| = ds \end{aligned}$$

So we take

$$\sigma = \frac{\lambda}{|f_z|} \circ f^{-1}.$$

□

## 3.2 Quasiconformal groups

**Def 3.2.1.** Suppose  $G$  is a group of quasiconformal maps of  $\mathbb{S}^2$ , we say that  $G$  is a quasiconformal group if there exists  $k$  such that all  $f \in G$  is  $k$ -quasiconformal.

**Exam 3.2.1.** Group of Möbius transformations  $M$  is a quasiconformal group with  $k = 0$ . Similarly, we can consider the conjugate group  $fMf^{-1}$  by some quasiconformal  $f$ , it is also a quasiconformal group.

**Prop 3.2.1.**

$$\mu_{f \circ g}(z) = \frac{\mu_g(z) + \mu_f(g(z))\overline{\theta(z)}}{1 + \mu_g(z)\mu_f(g(z))\theta(z)}, \theta = \frac{g_z}{g_{\bar{z}}}.$$

We denote  $T_g(\mu_f(g)) = \mu_{f \circ g}(z)$ .

*Proof.* Let  $h = f \circ g$ .

By proposition 3.1.2,

$$\mu_f \circ g = \frac{g_z}{g_{\bar{z}}} \frac{\mu_h - \mu_g}{1 - \overline{\mu_g} \mu_h}$$

So we obtain

$$\mu_h(z) = \frac{\mu_f(g(z)) + \theta(z)\mu_g(z)}{\theta(z) + \mu_f(g(z))\overline{\mu_g(z)}} = \frac{\mu_g(z) + \mu_f(g(z))\overline{\theta(z)}}{1 + \mu_g(z)\mu_f(g(z))\theta(z)}.$$

□

We now want to prove that actually every group of quasiconformal maps must be conjugated to a group of Möbius transformations. And so we first show the following lemma.

**Lemma 3.2.1.** Given a compact set  $X \in \mathbb{D}$ , there exists a barycenter  $b(X) \in \mathbb{D}$  such that  $A(b(X)) = b(A(X))$  for any  $A \in \text{Aut}(\mathbb{D})$ . More explicitly,  $b(X)$  can be given by the unique point such that

$$\int_X \frac{z - b(X)}{1 - \overline{b(X)}z} \frac{i}{2} dz \wedge d\bar{z} = 0.$$

To proof this lemma, we need some properties from analytic.

**Prop 3.2.2.** *The function*

$$G_X(z) = - \int_X \log \frac{(1 - |z|^2)(1 - |w|^2)}{|z - w|^2 + (1 - |z|^2)(1 - |w|^2)} \frac{i}{2} dw \wedge d\bar{w}$$

has a unique minimum in  $\mathbb{D}$ .

*Proof.*

$$\frac{(1 - |z|^2)(1 - |w|^2)}{|z - w|^2 + (1 - |z|^2)(1 - |w|^2)} = 1 - \frac{|z - w|^2}{|1 - \bar{z}w|^2} = 1 - \tanh^2 d(z, w) = \operatorname{sech}^2 d(z, w).$$

$$G_X(z) = \int_X \log \cosh^2 d(z, w) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2$$

Notice that  $d(z, w)$  is convex function of  $z$  for fixed  $w$  since  $\mathbb{D}$  is hyperbolic.

And  $\frac{d^2}{dt^2} \log \cosh^2 t = \operatorname{sech}^2 t > 0$ .

So  $G_X$  is convex.

Since  $G_X(z)$  tends to  $+\infty$  as  $|z| \rightarrow 1$ .

Hence  $G$  has a unique minimum in  $\mathbb{D}$ . □

*Proof of lemma 3.2.1.* By proposition 3.2.2, we take the minimum  $b \in \mathbb{D}$  of  $G_X$  and let

$$T_b(z) = \frac{z - b}{1 - \bar{b}z}, G(z) = G_X(T_b(z)).$$

Then we have

$$\begin{aligned} G(z) &= \int_A \log \cosh^2(d(T_b(z), w)) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2 \\ &= \int_A \log \cosh^2(d(z, T_b(w))) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2 \end{aligned}$$

So

$$\nabla G(z) = \int_A \left( \frac{2z}{1 - |z|^2} + \frac{2z|T_b(w)|^2 - 2T_b(w)}{|z - w|^2 + (1 - |z|^2)(1 - |w|^2)} \right) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2$$

In particular,  $\nabla G(0) = 0$ , *i.e.*

$$-2 \int_X T_b(w) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2 = 0$$

since  $z = 0$  is the minimum of  $G$ .

Take  $b(X) = b$ , we now prove that it is conformally invariant.

$$\begin{aligned} \int_{A(X)} T_{A(b(X))}(\zeta) \frac{i}{2} d\zeta \wedge d\bar{\zeta} / (1 - |\zeta|^2)^2 &= \int_X T_{A(b(X))}(A(w)) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2 \\ &= \int_X e^{i\theta} T_b(X)(w) \frac{i}{2} dw \wedge d\bar{w} / (1 - |w|^2)^2 \\ &= 0 \end{aligned}$$

where  $\zeta = A(w)$  and  $\theta \in [0, 2\pi)$ .

Hence  $A(b(X))$  is the unique minimum of  $G_{A(X)}(z)$ , *i.e.*  $A(b(X)) = b(A(X))$ . □

**Thm 3.2.1.** *Every quasiconformal group  $G$  is conjugated (by a quasiconformal map) to a subgroup of  $\operatorname{Aut}(\mathbb{S}^2)$ .*

*Proof.* Consider the sets  $M_z = \{\mu_f(z) | f \in G\}$  for  $z \in \mathbb{S}^2$ .

Then for  $g \in G$ ,  $T_g(M_{g(z)}) = \{T_g(\mu_f(g(z))) | f \in G\} = \{\mu_{f \circ g}(z) | f \in G\} = M_z$ .

Let  $\mu(z) = P(M_z)$  and solve Beltrami equation

$$\frac{h_{\bar{z}}}{h_z} = \mu.$$

Then we have

$$\begin{aligned} \mu_{h \circ g}(z) &= T_g(\mu_h(g(z))) = T_g(P(M_{g(z)})) \\ &= P(T_g(M_{g(z)})) = P(M_z) = \mu_h(z) \end{aligned}$$

So  $h \circ g \circ h^{-1}$  is conformal for every  $g \in G$ .  $\square$

In the proof we actually use the fact that the Beltrami equation can be solved, this will be proved later.

### 3.3 Holomorphic motions

**Def 3.3.1.** Let  $X$  be a connected complex manifold and  $E$  be a subset of  $\bar{\mathbb{C}}$ , we say a map  $f : E \times X \rightarrow \mathbb{S}^2$  is a holomorphic motion of  $E$  over  $X$  if

- (1) For fixed  $\lambda \in X$ ,  $f(-, \lambda) : E \rightarrow \mathbb{S}^2$  homeomorphic to its image.
- (2) For fixed  $z \in E$ ,  $f(z, -) : X \rightarrow \mathbb{S}^2$  is holomorphic.
- (3) For any  $z \in E$ ,  $f(z, \lambda_0) = z$  for some  $\lambda_0$ .

**Thm 3.3.1** (Schottky). *If  $h$  is analytic on  $\mathbb{D}$  and not equal to 0 or 1, then*

$$|h(z)| \leq \Phi \left( |h(0)|, \frac{1 + |z|}{1 - |z|} \right)$$

for  $|z| < 1$  where  $\Phi(x, y)$  is a universal strictly increasing continuous function for  $x \geq 0$  with  $\Phi(0, y) = 0$ .

**Lemma 3.3.1** ( $\lambda$ -lemma). *A holomorphic motion  $f(z, \lambda)$  of  $E = \mathbb{D}$  over  $\mathbb{D}$  can be extended to a quasiconformal map from  $\mathbb{D}$  into a closed quasidisk in  $\mathbb{C}$  in the  $z$  variable, and  $f(z, -)$  is holomorphic for any  $z \in \bar{\mathbb{D}}$ .*

*Proof.* Let  $z_1, z_2, z_3$  be three points in  $\mathbb{D}$  and

$$h(\lambda) = \frac{f(z_1, \lambda) - f(z_2, \lambda)}{f(z_3, \lambda) - f(z_2, \lambda)}.$$

WLOG, we assume  $\lambda_0 = 0$ .

Then by Schottky theorem,

$$\left| \frac{f(z_1, \lambda) - f(z_2, \lambda)}{f(z_3, \lambda) - f(z_2, \lambda)} \right| \leq \Phi \left( \left| \frac{z_1 - z_2}{z_3 - z_2} \right|, \frac{1 + |\lambda|}{1 - |\lambda|} \right).$$

Fixing  $z_2, z_3$ ,  $f(t, z)$  is bounded as  $z \in \mathbb{D}$  for a fixed  $\lambda$ .

And as  $z_1 \rightarrow z_2$ ,  $f(z_1, \lambda)$  tends to  $f(z_2, \lambda)$  since  $\Phi(0, \lambda) = 0$ .

So  $f(-, \lambda)$  is uniformly continuous, *i.e.* it can be continuously extended to  $\bar{\mathbb{D}}$ .

And by definition,  $f(-, \lambda)$  is quasisymmetric on  $\partial\mathbb{D}$ , *i.e.*  $f(-, \lambda)$  is quasiconformal on  $\bar{\mathbb{D}}$ .

On the other hand,  $f(z, -)$  is the uniform limit of some holomorphic maps for  $z \in \partial\mathbb{D}$ .

Hence  $f(z, -)$  is holomorphic for any  $z \in \bar{\mathbb{D}}$ .  $\square$

**Thm 3.3.2.** Suppose  $f(z, \lambda) : \mathbb{S}^2 \times \mathbb{D} \rightarrow \mathbb{S}^2$  is a holomorphic motion, then  $f(-, \lambda)$  is  $\frac{1+|\lambda|}{1-|\lambda|}$ -quasiconformal.

*Proof.* We restrict  $f$  on  $B_r \times \mathbb{D}$  and denote it as  $f_r$ , where  $B_r = \{z \mid |z| \leq R\}$ .

By  $\lambda$ -lemma,  $f_r(-, \lambda)$  is  $k$ -quasiconformal where  $k$  does not depend on  $r$  by lemma 2.2.1.

So as  $r \rightarrow \infty$ , we obtain that  $f(-, \lambda)$  is  $k$ -quasiconformal.

Let  $\mu(z, \lambda)$  be the Beltrami differential of  $f(-, \lambda)$ .

Then  $\mu(z, -)$  is holomorphic for any  $z \in \mathbb{S}^2$ .

So by Schwarz lemma,  $|\mu(z, \lambda)| \leq |\lambda|$ .

Hence  $f(-, \lambda)$  is  $\frac{1+|\lambda|}{1-|\lambda|}$ -quasiconformal. □

**Def 3.3.2.** Cross ratio of  $z_1, z_2, z_3, z_4 \in \mathbb{S}^2$  is

$$\text{Cr}(z_1, z_2, z_3, z_4) = \frac{z_4 - z_1}{z_4 - z_3} \cdot \frac{z_2 - z_3}{z_2 - z_1}.$$

**Prop 3.3.1.** Let  $f$  be a  $k$ -quasiconformal map, giving a compact set  $S \subset \mathbb{S}^2 \setminus \{0, 1, \infty\}$ , then

$$\{\text{Cr}(f(z_1), f(z_2), f(z_3), f(z_4)) \mid \text{Cr}(z_1, z_2, z_3, z_4) \in S\}$$

is compact.

*Proof.* Suppose the conclusion is wrong.

Then there exists  $(z_1^n, z_2^n, z_3^n, z_4^n)$  such that  $\text{Cr}(z_1^n, z_2^n, z_3^n, z_4^n) \in E$  but

$$\text{Cr}(f(z_1^n), f(z_2^n), f(z_3^n), f(z_4^n)) \rightarrow x \in \{0, 1, \infty\}.$$

Let  $A_n$  be the Möbius transformation that maps  $z_1^n, z_2^n, z_3^n$  to  $0, 1, \infty$  resp.

Then  $\text{Cr}(z_1^n, z_2^n, z_3^n, z_4^n) = A_n(z_4^n)$ .

WLOG, we assume  $A_n(z_4^n) \rightarrow z \in E$  by passing to a subsequence.

Similarly, consider  $B_n$  that maps  $f(z_1^n), f(z_2^n), f(z_3^n)$  to  $0, 1, \infty$  resp.

Then  $\text{Cr}(f(z_1^n), f(z_2^n), f(z_3^n), f(z_4^n)) = B_n(f(z_4^n))$ .

Let  $g_n = B_n \circ f \circ A_n^{-1}$ .

Then  $g_n$  fixes  $0, 1$  and  $\infty$ .

By lemma 1.7.1, we assume  $g_n \rightarrow g$  uniformly by passing to subsequence.

So  $\lim_{n \rightarrow \infty} g_n(A_n(z_4^n)) = g(z) = x \in \{0, 1, \infty\}$ , contradiction! □

Conversely, we can easily prove that:

**Thm 3.3.3.** There exists a compact set  $S \subset \mathbb{S}^2 \setminus \{0, 1, \infty\}$  such that if  $f : \mathbb{C} \rightarrow \mathbb{C}$  is a homeomorphism and  $N_f(S)$  is compact in  $\hat{\mathbb{C}} \setminus \{0, 1, \infty\}$ , then  $f$  is quasiconformal.

*Proof.* Take  $S = \{z \mid |z| = 2\}$ .

For  $z \in \mathbb{C}$ , we take  $x, y \in \partial B_r(z)$ , then

$$\frac{d(f(y), f(z))}{d(f(x), f(z))} = \frac{d(f(y), f(z))}{d(f(2y), f(z))} \frac{d(f(2y), f(z))}{d(f(x), f(z))}$$

Since  $\text{Cr}(z, x, \infty, 2y), \text{Cr}(z, y, \infty, 2y) \in S$ .

So similar to theorem 1.7.1,  $f$  is quasiconformal. □

Moreover, we consider a stronger proposition:

**Prop 3.3.2.** Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a homeomorphism, if for a fixed point  $z_0 \in \hat{\mathbb{C}} \setminus \{0, 1, \infty\}$ ,  $N_f(z_0)$  is compact in  $\hat{\mathbb{C}} \setminus \{0, 1, \infty\}$ , then  $f$  is quasiconformal.

This proposition is true but very difficult to prove. The main gap is that we can only estimate the dilatation for some specific angle ( $n\arg(z_0)$  for integer  $n$ ). In particular, when  $z_0$  is real, the condition seems to be very “weak”. I used to think it is wrong, but a intuitive understanding of why it is true is that every line (or circle) maps to  $\mathbb{C}$ , under  $f$ , quasimetrically with the same module  $K$ .

**Thm 3.3.4** (Słodkowski). *Every holomorphic motion  $f : E \times \mathbb{D} \rightarrow \mathbb{S}^2$  can be extended to holomorphic motion  $f : \mathbb{S}^2 \times \mathbb{D} \rightarrow \mathbb{S}^2$ .*

### 3.4 Two integral operators

We now start to solve the Beltrami equation.

**Def 3.4.1.** Let  $h \in L^p(\mathbb{C})$  with  $p > 2$ , the Cauchy operator is

$$Ph(\zeta) = -\frac{1}{\pi} \int_{\mathbb{C}} h(z) \left( \frac{1}{z-\zeta} - \frac{1}{z} \right) dx dy.$$

**Lemma 3.4.1.** *Ph is a continuous function with is Hölder with exponent  $1 - \frac{2}{p}$ .*

*Proof.* By Hölder inequality,

$$\begin{aligned} |Ph(\zeta)| &= \left| \frac{\zeta}{\pi} \int_{\mathbb{C}} h(z) \cdot \frac{1}{z(z-\zeta)} dx dy \right| \\ &\leq \frac{|\zeta|}{\pi} \|h\|_p \left\| \frac{1}{z(z-\zeta)} \right\|_q \end{aligned}$$

And since

$$\int_{\mathbb{C}} |z(z-\zeta)|^{-q} dx dy = |\zeta|^{2-2q} \int_{\mathbb{C}} |z(z-1)|^{-q} dx dy$$

So

$$|Ph(\zeta)| \leq \frac{1}{\pi} \left\| \frac{1}{z(z-1)} \right\|_q \|h\|_p |\zeta|^{1-\frac{2}{p}}.$$

Set  $h_1(z) = h(z + \zeta_1)$ , then

$$\begin{aligned} |Ph(\zeta_2) - Ph(\zeta_1)| &= |Ph_1(\zeta_2 - \zeta_1)| \\ &\leq K_p \|h\|_p |\zeta_1 - \zeta_2|^{1-\frac{2}{p}} \end{aligned}$$

□

**Def 3.4.2.** For  $h \in C_0^2(\mathbb{C})$ , we define

$$Th(\zeta) = \lim_{\varepsilon \rightarrow 0} -\frac{1}{\pi} \int_{|z-\zeta|>\varepsilon} \frac{h(z)}{(z-\zeta)^2} dx dy$$

**Lemma 3.4.2.** *If  $h \in C_0^2$ , then Th is  $C^1$  and*

$$(Ph)_z = Th, (Ph)_{\bar{z}} = h, \int |Th|^2 dx dy = \int |h|^2 dx dy.$$

*Proof.* Notice that

$$\begin{aligned}
(Ph)_{\bar{\zeta}} &= -\frac{1}{\pi} \int \frac{h_{\bar{z}}}{z-\zeta} dx dy \\
&= \frac{1}{2\pi i} \int \frac{h_{\bar{z}}}{z-\zeta} dz d\bar{z} \\
&= \lim_{\varepsilon \rightarrow 0} \frac{1}{2\pi i} \int_{|z-\zeta| > \varepsilon} \frac{h_{\bar{z}}}{z-\zeta} dz d\bar{z} \\
&= \lim_{\varepsilon \rightarrow 0} \frac{1}{2\pi i} \int_{|z-\zeta| = \varepsilon} \frac{h dz}{z-\zeta} = h(\zeta) \\
\int |Th|^2 dx dy &= -\frac{1}{2i} \int (Ph)_z \overline{(Ph)_z} dz d\bar{z} \\
&= \frac{1}{2i} \int Ph \overline{(\overline{Ph})_{\bar{z}z}} dz d\bar{z} \\
&= \frac{1}{2i} \int Ph \bar{h}_{\bar{z}} dz d\bar{z} \\
&= -\frac{1}{2i} \int (Ph)_{\bar{z}} |h|^2 \\
&= \int |h|^2 dx dy
\end{aligned}$$

□

*Remark 3.4.1.* This implies that  $Ph$  actually gives an solution of the  $\bar{\partial}$  problem.

**Thm 3.4.1** (Calderon-Zygmund). *Let  $h \in L^p(\mathbb{C})$  with  $p > 1$ , then  $\|Th\|_p \leq C_p \|h\|_p$  where  $c_p \rightarrow 1$  as  $p \rightarrow 2$ .*

**Conj 3.4.1.**  $\|T\|_p = p^* - 1$  where  $p^* = \max\left\{p, \frac{p}{p-1}\right\}$ .

*Remark 3.4.2.* The inequality  $\|T\|_p \geq p^* - 1$  is known.

### 3.5 Beltrami equations

**Thm 3.5.1.** *Suppose  $\mu$  has compact support and  $\|\mu\|_\infty \leq k < 1$ , fix  $p > 2$  such that  $kC_p < 1$ , then there exists a unique solution of  $f_{\bar{z}} = \mu f_z$  such that  $f(0) = 0$  and  $f_z - 1 \in L^p(\mathbb{C})$ .*

*Proof.* We first proof the uniqueness.

Since  $f_{\bar{z}} = f_z \mu$ .

So  $f_{\bar{z}} \in L^p$  and  $P(f_{\bar{z}})$  is well-defined.

Take  $F = f - P(f_{\bar{z}})$ .

Then  $F_{\bar{z}} = 0$  a.e., i.e.  $F$  is holomorphic.

Since  $F' = F_z = f_z - T(f_{\bar{z}})$ .

Therefore  $F' - 1 \in L^p(\mathbb{C})$ , i.e.  $F' - 1 = 0$ .

Thus  $f(z) = z + P(f_{\bar{z}})$ .

Suppose  $f, g$  is the solution of  $f_{\bar{z}} = \mu f_z$ , then

$$f_z - g_z = T(f_{\bar{z}} - g_{\bar{z}}) = T(\mu(f_z - g_z))$$

So by Calderon-Zygmund,

$$\|f_z - g_z\|_p \leq c_p \|\mu(f_z - g_z)\| \leq kc_p \|f_z - g_z\|,$$

contradiction!

Define  $h = T\mu + T\mu T\mu + \dots$ .

Since linear operator  $h \mapsto T(\mu h)$  has norm  $kC_q < 1$ .

So  $h$  is well-defined.

Take  $f = P(\mu(h+1)) + z$ .

Then  $f_z = T(\mu(h+1)) + 1 = h+1, f_{\bar{z}} = \mu(h+1)$ . □

**Coro 3.5.1.** *Suppose  $\mu_k \rightarrow \mu$  pointwisely a.e.,  $\|\mu_k\|, \|\mu\| < k$  and  $\text{supp } \mu_k, \text{supp } \mu \subset B(R)$ , then  $f_n \rightrightarrows f$ .*

*Proof.*

$$\begin{aligned} \|(f_n)_z - f_z\|_p &\leq \|T\mu((f_n)_z - f_z)\|_p + \|T(\mu - \mu_n)(f_n)_z\|_p \\ &\leq kC_p\|(f_n)_z - f_z\| + C_p\|(\mu_n - \mu)f_z\|_p \end{aligned}$$

As  $n \rightarrow \infty$ ,  $\|\mu_n - \mu\|$  tends to 0.

So  $\|(f_n)_z - f_z\|_p \rightarrow 0$ , i.e.  $f_n \rightrightarrows f$ . □

**Lemma 3.5.1.** *If  $\mu$  is smooth with compact support, then  $f$  is a topological mapping.*

*Proof.* Let  $f_z = \lambda$  and so  $\mu\lambda = f_{\bar{z}}$ , then

$$\lambda_{\bar{z}} = (\mu\lambda)_z = \lambda_z\mu + \lambda\mu_z.$$

$$(\log \lambda)_{\bar{z}} = \mu(\log \lambda)_z + \mu_z.$$

Take  $\sigma = \log \lambda$ , then

$$\sigma_{\bar{z}} = \mu\sigma_z + \mu_z.$$

Let  $q \in L^p(\mathbb{C})$  such that

$$q = T(\mu q) + T\mu_z.$$

If such  $q$  exists, then we can let

$$\sigma = P(\mu q + \mu_z) + C.$$

To find  $q$ , we consider operator  $T : L^p \rightarrow L^p, q \mapsto T(\mu q) + T\mu_z$ .

By Schauder fixed point theorem,  $q$  exists.

So  $|f_z| = |e^\sigma| > 0$ , i.e.  $f$  is a topological mapping. □

**Thm 3.5.2.** *If  $\mu$  with compact support and  $\|\mu\|_\infty \leq k < 1$ , then the solution of Beltrami equation is quasiconformal homeomorphism.*

**Thm 3.5.3.** *For any measurable function  $\mu$  on  $\mathbb{C}$  with  $\|\mu\|_\infty \leq k < 1$ , there exists a unique quasiconformal map  $f^\mu$  such that  $f_z^\mu = \mu f_z^\mu$  a.e., and  $f^\mu(0) = 0, f^\mu(1) = 1, f^\mu(\infty) = \infty$ .*

*Proof.* We first consider the special case that  $\mu = 0$  near 0.

Then take

$$\hat{\mu}(z) = \mu \left( \frac{1}{z} \right) \frac{z^2}{\bar{z}^2}.$$

So  $\hat{\mu}$  has compact support and

$$f^\mu(z) = \frac{1}{f^{\hat{\mu}}\left(\frac{1}{z}\right)}.$$

For general case, take  $\mu = \mu_1 + \mu_2$  where  $\mu_1$  has compact support and  $\mu_2 = 0$  near 0.

Take

$$\lambda = \left( \frac{\mu - \mu_2}{1 - \mu\bar{\mu}_2} \frac{f_z^{\mu_2}}{f_z^{\mu_2}} \right) \circ (f^{\mu_2})^{-1}.$$

Then  $\lambda$  has compact support and

$$f^\mu = f^\lambda \circ f^{\mu_2}.$$

□

**Thm 3.5.4.** *Suppose  $X$  is a Riemann surface, and let  $\mu$  be a  $(-1, 1)$  form on  $X$  with  $\|\mu\|_\infty < 1$ , then there exists a new Riemann surface  $X^\mu$  and a quasiconformal map  $f^\mu : X \rightarrow X^\mu$  such that*

$$\bar{\partial}f^\mu = \mu\partial f^\mu.$$

*Proof.* Let  $X = \mathbb{H}/\Gamma$  and lift  $\mu$  to  $\tilde{\mu}$  on  $\mathbb{H}$  satisfying that

$$(\tilde{\mu} \circ A) \frac{\overline{A'}}{A'} = \tilde{\mu},$$

for any  $A \in \Gamma$ .

Extend  $\tilde{\mu}$  to  $\mathbb{C}$  by taking  $\overline{\tilde{\mu}(\bar{z})}$  under the real-axis and take  $f = f^{\tilde{\mu}}$ , then

$$\begin{aligned} \text{Belt}(f \circ A) &= (\text{Belt}(f) \circ A) \frac{\overline{A'}}{A'} \\ &= (\tilde{\mu} \circ A) \frac{\overline{A'}}{A'} \\ &= \tilde{\mu} = \text{Belt}(f) \end{aligned}$$

So  $f \circ A = B \circ f$  for some Möbius transformation  $B$ .

Hence  $\Gamma_\mu = f \circ \Gamma \circ f^{-1}$  is a Fuchsian group and we can take  $X^\mu = \mathbb{H}/\Gamma_\mu$ .  $\square$

*Remark 3.5.1.* We can extend  $\mu$  arbitrarily and use Riemann mapping theorem to let  $f$  maps  $\mathbb{D}$  to  $\mathbb{D}$ .

**Exam 3.5.1.** *Take  $f(x, y) = (2x, y)$  in  $0 \leq x \leq 1$  and  $f(x, y) = (\frac{1}{2}x + \frac{3}{2}, y)$ .*

*Then we extend  $f$  to  $\mathbb{C}$  by  $f(x + 3n, y) = f(x, y) + 3n$  and so*

$$\text{Belt}(f)(x, y) = \begin{cases} \frac{1}{3} & 3n \leq x \leq 3n + 1 \\ -\frac{1}{3} & 3n + 1 \leq x \leq 3n + 3 \end{cases}$$

*Let  $f_n(z) = \frac{1}{n}f(nz)$ .*

*Then  $f_n(z) \rightarrow z$  as  $n \rightarrow \infty$ , but  $\mu_n \not\rightarrow 0$  in distribution sense.*

*Moreover,  $\mu_n \rightarrow -\frac{1}{9}$  in weak\* topology.*

## 3.6 Decomposition of quasiconformal maps

**Thm 3.6.1.** *Let  $f$  be a  $k$ -quasiconformal map and  $0 < t < 1$ , then  $f = f_2 \circ f_1$  where  $f_1$  is  $K^t$ -quasiconformal and  $f_2$  is  $K^{1-t}$ -quasiconformal.*

*Proof.* Let  $\frac{\mu_1(z)}{\mu(z)} \in \mathbb{C}$  with  $d(0, \mu_1(z)) = td(0, \mu(z))$ , then

$$\frac{1 + |\mu_1(z)|}{1 - |\mu_1(z)|} = \left( \frac{1 + |\mu(z)|}{1 - |\mu(z)|} \right)^t.$$

So  $f_1 = f^{\mu_1}$  is  $K^t$ -quasiconformal.

Take  $f_2 = f \circ f_1^{-1}$  and  $\mu_2 = \text{Belt}(f_2)$ , then

$$|\mu_2(f_1(z))| = \left| \frac{\mu - \mu_1}{1 - \bar{\mu}_1\mu} \right|$$

Therefore we obtain

$$\log \left( \frac{1 + |\mu_2(f_1(z))|}{1 - |\mu_2(f_1(z))|} \right) = 2d(\mu_1(z), \mu(z)) = (1 - t) \log \left( \frac{1 + |\mu(z)|}{1 - |\mu(z)|} \right).$$

Hence  $f_2$  is  $K^{1-t}$  quasiconformal.  $\square$

**Coro 3.6.1.** *Let  $\varepsilon > 0$  and  $f$  be a  $k$ -quasiconformal map, then  $f = f_1 \circ \dots \circ f_n$  where  $(1 + \varepsilon)$ -quasiconformal maps  $f_1, \dots, f_n$  for sufficiently large  $n$ .*

*Proof.* Take  $n$  such that  $(1 + \varepsilon)^n > k$  and  $g_1 = f$ .

By the above theorem, we construct  $g_i = f_i \circ g_{i+1}$  where  $f_i$  is  $K^{\frac{1}{n}}$ -quasiconformal inductively.  $\square$

**Conj 3.6.1.** *Let  $\varepsilon > 0$  and  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a  $k$ -quasiconformal map, are there  $(1 + \varepsilon)$ -quasiconformal maps  $f_1, \dots, f_k$ , such that  $f = f_1 \circ \dots \circ f_k$ .*

**Conj 3.6.2.** *Let  $\varepsilon > 0$  and  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a  $L$ -biLipschitz map, are there  $f_1, \dots, f_r$  which are  $(1 + \varepsilon)$ -biLipschitz such that  $f = f_1 \circ \dots \circ f_r$ .*

### 3.7 Dependence on parameter

**Def 3.7.1.** Let  $\nu$  be an essentially bounded measurable function in  $\mathbb{C}$ , for small  $t > 0$  with  $\|t\nu\|_\infty < 1$ , we denote

$$f^{t\nu}(z) = z + t\dot{f}[\nu](z) + o(t), \text{ i.e. } \dot{f}[\nu](z) = \lim_{t \rightarrow 0} \frac{f^{t\nu}(z) - z}{t}$$

**Prop 3.7.1.** *Suppose  $\nu$  is compact support and Let  $f^{t\nu}$  be the unique solution such that  $f(0) = 0$  and  $f_z - 1 \in L^p$ , then*

$$\dot{f}[\nu](z) = -\frac{1}{\pi} \int_{\mathbb{C}} \nu(w) \left( \frac{1}{w-z} - \frac{1}{w} \right) dudv$$

*Proof.*

$$f^{t\nu} = z + P(t\nu + t\nu T(t\nu) + \dots).$$

So we obtain

$$\dot{f}[\nu](z) = P(\nu) = -\frac{1}{\pi} \int_{\mathbb{C}} \nu(w) \left( \frac{1}{w-z} - \frac{1}{w} \right) dudv$$

$\square$

**Thm 3.7.1.** *Let  $f^{t\nu}$  be the unique solution fixing  $0, 1$  and  $\infty$ , which we call the normalized quasiconformal map, then*

$$\dot{f}[\nu](z) = -\frac{1}{\pi} \int_{\mathbb{C}} \nu(w) \frac{z(z-1)}{w(w-1)(w-z)} dudv$$

**Coro 3.7.1.** *Let  $\{\mu(t)\}$  be an family of Beltrami coefficients depending on a real or complex parameter  $t$  such that  $\mu(t)$  is differentiable at  $t = 0$ , that is,*

$$\mu(t)(z) = \mu(z) + t\nu(z) + t\varepsilon(t)(z)$$

for  $z \in \mathbb{C}$  and  $\nu, \varepsilon(t) \in L^\infty(\mathbb{C})$  such that  $\|\varepsilon(t)\|_\infty \rightarrow 0$  as  $t \rightarrow 0$ . Then for normalized quasiconformal maps  $f^{\mu(t)}$ ,

$$\dot{f}[\nu](z) = -\frac{1}{\pi} \int_{\mathbb{C}} \nu(w) \frac{f^\mu(z)(f^\mu(z) - 1) ((f^\mu)_z(w))^2}{f^\mu(w)(f^\mu(w) - 1)(f^\mu(w) - f^\mu(z))} dudv$$

# Chapter 4

## Teichmüller space

### 4.1 Integrable holomorphic quadratic differential

**Def 4.1.1.**  $X$  is a Riemann surface, a holomorphic quadratic differential  $\phi$  on  $X$  is a holomorphic  $(2,0)$ -form.

**Prop 4.1.1.** Let  $\phi = \phi(z)dz^2$  be a holomorphic quadratic differential, then  $|\phi|$  is a volume form on  $X$ .

*Proof.* Under holomorphic change of coordinates,  $\phi(A(w))A'(w)^2 = \tilde{\phi}(w)$ . □

**Def 4.1.2.** Denote the integrable quadratic differentials by  $\text{MQD}^1(X)$  and integrable holomorphic quadratic differentials by  $\text{QD}^1(X)$ .

We define  $B^\infty(X) = \{\mu | \mu \text{ is } (-1, 1) \text{ measurable form with } \|\mu\|_\infty < +\infty\}$ .

**Prop 4.1.2.** Take  $\phi \in \text{QD}^1(X), \mu \in B^\infty(X)$ , then  $\phi\mu$  is a volume form.

*Proof.* Under holomorphic change of coordinates,

$$\tilde{\phi}(w) = (\phi\mu)(A(w)) \frac{\overline{A'(w)}}{A'(w)} A'(w)^2 = (\phi\mu)(A(w)) A'(w) \overline{A'(w)}$$

□

**Prop 4.1.3.** Suppose  $\lambda : \text{QD}^1(X) \rightarrow \mathbb{C}$  is a bounded linear function, then it can extend to  $\lambda : \text{MQD}^1(X) \rightarrow \mathbb{C}$  with the same norm and there exists  $\mu \in B^\infty(X)$  such that for any  $\phi \in \text{QD}^1(X)$ , we have

$$\lambda(\phi) = \int_X \mu\phi, \|\mu\|_\infty = \|\lambda\|.$$

Moreover,  $\mu$  is unique if  $\dim \text{QD}^1(X) < \infty$ .

*Proof.* By Hahn-Banach theorem, we can extend  $\lambda$  to  $\lambda : \text{MQD}^1(X) \rightarrow \mathbb{C}$  with the same norm. And by Riesz representation theorem, such  $\mu$  exists.

For the uniqueness, suppose  $\|\lambda\| = 1$  and  $\dim \text{QD}^1(X) < \infty$ .

Then there exists  $\phi_0 \in \text{QD}^1(X)$  such that  $|\lambda(\phi_0)| = \|\phi_0\|_1$ , so

$$\|\phi_0\|_1 = \int_x \mu\phi_0 \leq \|\mu\|_\infty \|\phi_0\|_1.$$

Hence the only possibility is that  $\mu = \frac{|\phi_0|}{\phi_0}$ . □

**Exam 4.1.1.** Suppose  $X = \{z | 0 < \text{Im } z < 1\}$  is the strip,  $\lambda(\phi) = \int_X \phi$  with  $\phi dz^2 \in \text{QD}^1(X)$ .

Take  $\phi_n = \frac{1}{n+z^2}$ , then

$$\int_X \phi_n = \int_{-\infty}^{\infty} \frac{dx}{n+x^2}, \int_X |\phi_n| = \int_X \frac{1}{n \left| 1 + \frac{z^2}{n} \right|} = \frac{\sqrt{n}}{n} \int_0^1 dy \int_{-\infty}^{\infty} \frac{dx}{\left| 1 + \left( x + \frac{y}{\sqrt{n}} \right)^2 \right|}$$

So

$$\lim_{n \rightarrow \infty} \frac{|\lambda(\phi_n)|}{\|\phi_n\|_1} = 1.$$

Hence  $\|\lambda\| = 1$ .

**Exam 4.1.2.** Suppose  $X = \mathbb{H}$ ,  $\lambda(\phi) = \int_X \phi$  with  $\phi dz^2 \in \text{QD}^1(X)$ .

Then by Cauchy theorem,  $\lambda(\phi) = 0$  since  $\phi$  is integrable.

**Lemma 4.1.1.** Meromorphic  $(2, 0)$ -forms with first order poles on  $E = \partial\mathbb{D}$  are dense in  $\text{QD}^1(\mathbb{D})$

*Proof.* Let  $L \subset \text{QD}^1(\mathbb{D})$  be the closure of such meromorphic functions.

Then we want to show  $L = \text{QD}^1(\mathbb{D})$ , suppose not.

So there exists a nonzero bounded linear functional  $\lambda : \text{QD}^1(\mathbb{D}) \rightarrow \mathbb{C}$  such that  $\lambda(L) = 0$ .

By proposition 4.1.3, we take such unique  $\mu \in B^\infty(\mathbb{D})$ .

Let

$$Q(z) = -\frac{1}{\pi} \int_{\mathbb{D}} \frac{\mu(w)}{w-z} dudv.$$

Then  $Q_{\bar{z}}(z) = \mu(z)$ ,  $Q(z) = 0$  for  $|z| = 1$  and  $Q$  is continuous on  $\mathbb{C}$ .

Suppose  $\phi dz^2 \in \text{QD}^1(\mathbb{D})$  is holomorphic in some neighborhood of  $\mathbb{D}$ , then

$$\int_{\mathbb{D}} \mu(z) \phi(z) dz d\bar{z} = \int_{\mathbb{D}} Q_{\bar{z}}(z) \phi(z) dz d\bar{z} = \int_{\partial\mathbb{D}} Q(z) \phi(z) dz = 0.$$

Now for any  $\phi dz^2 \in \text{QD}^1(\mathbb{D})$ , we let  $\phi_r(z) = \phi(rz)$  for  $r < 1$ , then

$$\int \mu(z) \phi_r(z) dz d\bar{z} = 0$$

So  $\int \mu \phi = 0$  since  $\phi_r \xrightarrow{L^1} \phi$  as  $r \rightarrow 1$ , contradiction! □

**Def 4.1.3.** Suppose  $X = \mathbb{D}/\Gamma$  is a Riemann surface, we define the Poincaré series operator as

$$\begin{aligned} \theta: \text{QD}^1(\mathbb{D}) &\longrightarrow \text{QD}^1(X) \\ \phi dz^2 &\longmapsto \sum_{A \in \Gamma} (\phi \circ A) (A')^2 dz^2 \end{aligned}$$

**Prop 4.1.4.**  $\theta : \text{QD}^1(\mathbb{D}) \rightarrow \text{QD}^1(X)$  is well-defined and  $\|\theta\| \leq 1$ .

*Proof.* For  $A \in \Gamma$  and  $w = A(z)$ ,

$$\begin{aligned} \theta(\phi dz^2)(w) &= \sum_{B \in \Gamma} \phi(B(w)) \cdot (B'(w))^2 dw^2 \\ &= \sum_{B \in \Gamma} \phi(B \circ A(z)) \cdot (B'(A(z)) \cdot A'(z)) dz^2 \\ &= \sum_{B \in \Gamma} \phi(B(z)) (B'(z))^2 dz^2 \end{aligned}$$

Moreover, let  $F$  be a fundamental domain of  $X$ , then

$$\begin{aligned}
 \|\theta(\phi dz^2)\|_1 &= \int_F \left| \sum_{A \in \Gamma} (\phi \circ A)(A')^2 \right| dx dy \\
 &\leq \sum_{A \in \Gamma} \int_F |\phi(A(z))| |A'(z)|^2 dx dy \\
 &= \sum_{A \in \Gamma} \int_{A(F)} |\phi(z)| dx dy \\
 &= \int_{\mathbb{D}} |\phi(z)| dx dy = \|\phi\|_1
 \end{aligned}$$

□

**Thm 4.1.1.** *Given  $\psi \in \text{QD}^1(X)$ , there exists  $\phi \in \text{QD}^1(\mathbb{D})$  such that  $\theta(\phi) = \psi$  and  $\|\phi\| \leq 3\|\psi\|$ .*

*Remark 4.1.1.* We will prove this in next section.

**Thm 4.1.2.** *Let  $\nu \in B^\infty(\mathbb{H}/\Gamma)$ , then*

(1)  $\dot{f}[\nu](z) = 0$  for any  $z \in \partial\mathbb{H}$  iff  $\int_X \nu \phi = \int_{\mathbb{H}/\Gamma} \nu \phi = 0$  for any  $\phi \in \text{QD}^1(X) \cong \text{QD}^1(\mathbb{H}/\Gamma)$ .

(2) Moreover, in this case there exists  $\delta(t) \in \text{Belt}(\mathbb{H}/\Gamma)$  such that

- a)  $\|\delta(t)\| \leq 12t^2$ .
- b)  $f^{t\nu}(z) = f^{\delta(t)}(z)$  for  $z \in \partial\mathbb{H}$ .

*Proof.* (1) By theorem 3.7.1

$$\dot{f}[\nu](z) = -\frac{1}{\pi} \int_{\mathbb{C}} \nu(w) \left( \frac{1}{w-z} - \frac{z}{w-1} - \frac{1}{w} \right)$$

So  $\left(\dot{f}[\nu]\right)_{\bar{z}} = \nu$ .

For meromorphic map  $\phi$  with simple pole in  $\partial\mathbb{H}$ ,

$$\begin{aligned}
 \int_{\mathbb{H}} \nu \phi dz dy &= \frac{1}{2i} \int_{\mathbb{H}} \left(\dot{f}[\nu]\right)_{\bar{z}} \phi dz d\bar{z} \\
 &= \frac{1}{2i} \int_{\partial\mathbb{H}} \dot{f}[\nu] \phi dz = 0
 \end{aligned}$$

By lemma 4.1.1,  $\int_{\mathbb{H}} \nu \phi = 0$  for any  $\phi \in \text{QD}^1(\mathbb{H})$ .

And since

$$\begin{aligned}
 \int_X \nu \theta(\phi dz^2) &= \int_F \nu \sum_{A \in \Gamma} (\phi \circ A)(A')^2 dz^2 \\
 &= \sum_{A \in \Gamma} \int_F \nu(\phi \circ A)(A')^2 dz^2 \\
 &= \sum_{A \in \Gamma} \int_{A(F)} (\nu \circ A^{-1}) \frac{\overline{A_z^{-1}}}{A_z} \phi dA(z)^2 \\
 &= \sum_{A \in \Gamma} \int_{A(F)} \nu \phi = \int_{\mathbb{H}} \nu \phi.
 \end{aligned}$$

Hence  $\int_X \nu\phi = 0$  for any  $\phi \in \text{QD}^1(X)$ .

The converse is trivial by taking

$$\phi(w) = \left( \frac{1}{w-z} - \frac{z}{w-1} - \frac{1}{w} \right) dz^2.$$

(2) It is too technical to prove, so we will not give the proof. □

## 4.2 Bergman projection

We denote the hyperbolic metric by  $\rho(z) = \frac{2}{1-|z|^2}$ .

**Def 4.2.1.** Let  $A_2(\mathbb{D})$  be the space of holomorphic functions  $\varphi$  on  $\mathbb{D}$  such that

$$\|\phi\|^2 = \int_{\mathbb{D}} \rho(z)^{-2} |\phi(z)|^2 dx dy < \infty.$$

We define the Petersson scalar product on  $A_2(\mathbb{D})$  by

$$\langle \phi, \psi \rangle_{\mathbb{D}} = \int_{\mathbb{D}} \rho(z)^{-2} \phi(z) \overline{\psi(z)} dx dy.$$

Then  $A_2(\mathbb{D})$  becomes a separable Hilbert space with this scalar product.

We take

$$\phi_n(z) = \sqrt{\frac{2}{\pi}} (n+1)(n+2)(n+3) z^n.$$

Then  $\{\phi_n\}_{n=0}^{\infty}$  is a complete orthonormal basis for  $A_2(\mathbb{D})$ .

**Def 4.2.2.** The Bergman kernel is the reproducing kernel for  $A_2(\mathbb{D})$ , that is,

$$K(z, w) = \sum_{n=0}^{\infty} \phi_n(z) \overline{\phi_n(w)} = \frac{12}{\pi(1-z\bar{w})^4}.$$

**Prop 4.2.1** (reproducing formula). For  $\phi \in A_2(\mathbb{D})$ ,

$$\phi(z) = \langle \phi, K(\cdot, z) \rangle_{\mathbb{D}}.$$

*Proof.*

$$\langle \phi, K(\cdot, z) \rangle_{\mathbb{D}} = \sum_{n=0}^{\infty} \langle \phi, \phi_n \cdot \overline{\phi_n(z)} \rangle_{\mathbb{D}} = \sum_{n=0}^{\infty} \phi_n(z) \langle \phi, \phi_n \rangle_{\mathbb{D}} = \phi(z)$$

□

**Thm 4.2.1.** (1)  $K(z, w) = \overline{K(w, z)}$

(2) For  $A \in \text{Aut}(\mathbb{D})$ ,  $K(Az, Aw) A'(z)^2 \overline{A'(w)^2} = K(z, w)$

(3)  $\int_{\mathbb{D}} |K(z, w)| dx dy = 3\rho^2(w)$

(4) If  $|\rho^{-2}(z)\phi(z)| \leq C$  for  $z \in \mathbb{D}$  and holomorphic  $\phi$ , then

$$\phi(z) = \int_{\mathbb{D}} \rho^{-2}(w) \phi(w) K(z, w) du dv$$

*Proof.* (1) Trivial.

(2) Since

$$\begin{aligned}
& \langle (\phi \circ A)(A')^2, (\psi \circ A)(A')^2 \rangle_{\mathbb{D}} \\
&= \int_{\mathbb{D}} \rho(z)^{-4} \phi(A(z)) \overline{\psi(A(z))} A'(z)^2 \overline{A'(z)}^2 dA_{\mathbb{D}}(z) \\
&= \int_{\mathbb{D}} (\rho(z)^{-1} |A'|)^4 \phi(A(z)) \overline{\psi(A(z))} dA_{\mathbb{D}}(A(z)) \\
&= \int_{A(\mathbb{D})} \rho(z)^{-4} \phi(z) \overline{\psi(z)} dA_{\mathbb{D}}(z) = \langle \phi, \psi \rangle_{\mathbb{D}}
\end{aligned}$$

Here

$$dA_{\mathbb{D}}(z) = \frac{4dx dy}{(1 - |z|^2)^2}$$

is the hyperbolic area.

So  $\{(\phi_n \circ A)(A')^2\}$  is also a complete orthonormal basis for  $A_2(\mathbb{D})$ .

Hence the reproducing kernel is the same, *i.e.*

$$K(Az, Aw) A'(z)^2 \overline{A'(w)}^2 = K(z, w)$$

(3)

$$\begin{aligned}
\int_{\mathbb{D}} |K(z, w)| dx dy &= \frac{12}{\pi} \int_0^1 \int_0^{2\pi} \frac{r}{(1 - 2sr \cos \theta + s^2 r^2)^2} d\theta dr \\
&= \frac{12}{\pi} \int_0^1 \frac{2\pi(1 + s^2 r^2)}{(1 - s^2 r^2)^3} dr \\
&= 24 \int_1^{1-s^2} \frac{2-t}{t^3} \cdot \frac{-dt}{2s^2} \\
&= \frac{12}{s^2} \int_{1-s^2}^1 \frac{2-t}{t^3} dt \\
&= \frac{12}{s^2} \left( \frac{1}{(1-s^2)^2} - \frac{1}{1-s^2} \right) \\
&= \frac{12}{(1-s^2)^2} = 3\rho^2(w)
\end{aligned}$$

Here  $z = re^{i\theta}$ ,  $w = se^{i\varphi}$ ,  $t = 1 - r^2 s^2$

(4) Since  $|\rho^{-2}(z)\phi(z)| \leq C$ .

So the integral is well-defined for all  $z$ .

For  $z = 0$ ,

$$\begin{aligned}
\phi(0) &= 6\phi(0) \int_0^1 (1-r^2)^2 r dr \\
&= \frac{3}{\pi} \int_{\mathbb{D}} (1-|w|^2)^2 \phi(w) du dv \\
&= \int_{\mathbb{D}} \rho(w)^{-2} \phi(w) K(0, w) du dv
\end{aligned}$$

Now for any  $z \in \mathbb{D}$ , take  $A \in \text{Aut}(\mathbb{D})$  such that  $A(0) = z$  and  $\psi = (\phi \circ A)(A')^2$ .

Then we obtain that

$$\begin{aligned}\psi(0) &= \int_{\mathbb{D}} \rho(\zeta)^{-2} \psi(\zeta) K(0, \zeta) d\xi d\eta \\ &= \int_{\mathbb{D}} (\rho(A(\zeta)) |A(\zeta)|)^{-2} \phi(A(\zeta)) A'(\zeta)^2 K(A(0), A(\zeta)) A'(0)^2 \overline{A'(\zeta)}^2 |A(\zeta)|^{-2} dudv \\ &= A'(0)^2 \int_{\mathbb{D}} \rho(w)^{-2} \phi(w) K(z, w) dudv\end{aligned}$$

Hence this conclude the desired formula. □

**Def 4.2.3** (Bergman projection).

$$P : MQD^1(\mathbb{D}) \rightarrow QD^1(\mathbb{D}), P(\phi)(z) = \int_{\mathbb{D}} \rho^{-2}(w) \phi(w) K(z, w) dudv.$$

*Proof of theorem 4.1.1.* Let  $F$  be a fundamental domain of  $X$  and take

$$\psi(z) = P(\chi_F \phi)(z) = \int_{\mathbb{H}} \rho^{-2}(w) (\chi_F(w) \phi(w)) K(z, w) dudv.$$

Then  $\theta\psi = \phi$  and

$$\begin{aligned}\int_{\mathbb{H}} |\psi(z)| dx dy &\leq \int_{\mathbb{H}} |\rho^{-2} \chi_F \phi|(w) \left( \int_{\mathbb{H}} |K(z, w)| dx dy \right) dudv \\ &\leq \int_{\mathbb{H}} |\rho^{-2} \chi_F \phi|(w) \cdot 3\rho^2(w) dudv \\ &= 3 \int_F |\phi(w)| dudv = 3\|\phi\|_1\end{aligned}$$

□

### 4.3 Teichmüller spaces

**Def 4.3.1.** Let  $S_0$  be a Riemann Surface, we let  $T(S_0) = \{(S, f) | f : S_0 \rightarrow S \text{ quasiconformal}\} / \sim$  and  $(S_1, f_1) \sim (S_2, f_2)$  if  $f_2 \circ f_1^{-1} : S_1 \rightarrow S_2$  is homotopic to a conformal map.

$\text{Belt}(S_0)$  is the unit ball of  $(-1, 1)$ -form on  $S_0$  w.r.t.  $\|\cdot\|_{\infty}$ .

*Remark 4.3.1.*  $T(S_0) = \text{Belt}(S_0) / \sim$  where  $\mu \sim \nu$  iff  $f^{\nu} \circ (f^{\mu})^{-1}$  is homotopic to conformal map.

**Def 4.3.2.**  $f_{\mu} : \mathbb{C} \rightarrow \mathbb{C}$  is the unique normalized quasiconformal map with Beltrami differential

$$\begin{cases} \mu(z) & z \in \mathbb{H} \\ 0 & z \in \mathbb{R} \cup \mathbb{H}^- \end{cases}$$

**Prop 4.3.1.**  $f^{\mu} = f^{\nu}$  on  $\mathbb{R}$  iff  $f_{\mu} = f_{\nu}$  on  $\mathbb{R}$ .

*Proof.* If  $f_{\mu} = f_{\nu}$  on  $\mathbb{R}$ , then  $f_{\mu} \circ (f^{\mu})^{-1}, f_{\nu} \circ (f^{\nu})^{-1}$  are conformal maps which map  $\mathbb{H}$  to  $\Omega$ , whose boundary is  $f_{\mu}(\mathbb{R})$ , and fix  $0, 1$  and  $\infty$ .

Then  $f_{\mu} \circ (f^{\nu})^{-1} = f_{\nu} \circ (f^{\nu})^{-1}$  on  $\overline{\mathbb{H}}$ .

And since  $f_{\mu} = f_{\nu}$  on  $\mathbb{R}$ .

So  $f^{\mu} = f^{\nu}$  on  $\mathbb{R}$ .

Conversely, we take

$$g(z) = \begin{cases} ((f^\nu)^{-1} \circ f^\mu)(z) & z \in \mathbb{H} \\ z & z \in \mathbb{R} \cup \mathbb{H}^- \end{cases}$$

And let  $A = f_\gamma \circ g \circ (f_\mu)^{-1}$ .

Then  $A$  is conformal on  $\Omega = f_\mu(\mathbb{H})$  since  $A = f_\gamma \circ (f^\gamma)^{-1} \circ f^\mu \circ (f_\mu)^{-1}$  on  $\Omega$ .

And similarly,  $A$  is conformal on  $\Omega^*$  and  $A$  is quasiconformal on  $\mathbb{C}$ .

So  $A$  is conformal on  $\mathbb{C}$  and fix  $0, 1$  and  $\infty$ , *i.e.*  $A = \text{Id}$ .

Hence  $f_\mu = f_\nu$  on  $\mathbb{R}$ . □

**Def 4.3.3.** We define the Teichmüller metric on Teichmüller space as

$$d((S_1, f_1), (S_2, f_2)) = \frac{1}{2} \inf_{g \sim f_2 \circ f_1^{-1}} \log K_g$$

**Lemma 4.3.1.** *Let  $f : S_0 \rightarrow S$  be quasiconformal, then the set of all quasiconformal maps  $g : S_0 \rightarrow S$  which are homotopic to  $f$  contains an extremal map with the smallest maximal dilatation.*

*Proof.* The lemma is trivial if the universal covering is  $\hat{\mathbb{C}}$  or  $\mathbb{C}$  and  $M$  is not compact.

The torus case is similar as below.

Now suppose  $\mathbb{H}$  is the universal covering of  $M$ .

Then lift  $g$  to a map  $\tilde{g} : \mathbb{H} \rightarrow \mathbb{H}$  such that all maps  $\tilde{g}$  agrees on boundary.

So the set of  $\tilde{g}$  contains its limits, *i.e.* there exists a map  $\tilde{g}_0$  with smallest maximal dilatation.

Hence  $g_0$  is the required extremal map. □

**Prop 4.3.2.** *Teichmüller metric is a metric.*

*Proof.* Take  $(S_1, f_1), (S_2, f_2), (S_3, f_3)$ , then

$$\begin{aligned} d([S_1, f_1], [S_2, f_2]) &= \frac{1}{2} \inf_{g \sim f_2 \circ f_1^{-1}} \log K_g \\ &\leq \frac{1}{2} \inf_{\substack{g_1 \sim f_2 \circ f_3^{-1} \\ g_2 \sim f_3 \circ f_1^{-1}}} \log K_{g_1} K_{g_2} \\ &= d([S_1, f_1], [S_3, f_3]) + d([S_2, f_2], [S_3, f_3]) \end{aligned}$$

Suppose  $d([S_1, f_1], [S_2, f_2]) = 0$ .

Then there exists extremal map  $g \sim f_2 \circ f_1^{-1}$  such that  $K_g = 1$ , *i.e.*  $g$  is conformal.

So  $[S_1, f_1] = [S_2, f_2]$ . □

*Remark 4.3.2.* In terms of Beltrami differentials,

$$d_T([\mu_0], [\nu_0]) = \frac{1}{2} \inf_{\mu \in [\mu_0], \nu \in [\nu_0]} \log \frac{1 + \left\| \frac{\mu - \nu}{1 - \bar{\mu}\nu} \right\|_\infty}{1 - \left\| \frac{\mu - \nu}{1 - \bar{\mu}\nu} \right\|_\infty}$$

**Def 4.3.4.** A Beltrami differential  $\mu \in [\mu_0]$  is called extremal if  $\|\mu\|_\infty \leq \|\nu\|_\infty$  for any  $\nu \in [\mu_0]$ .

**Prop 4.3.3.** *Let  $[\mu_1] \in T(S_0)$ , if  $\mu$  is an extremal Beltrami differential in the Teichmüller class  $[\mu_1]$ , then*

$$\mu_t = \frac{(1 + |\mu|)^t - (1 - |\mu|)^t}{(1 + |\mu|)^t + (1 - |\mu|)^t} \cdot \frac{\mu}{|\mu|}$$

for  $0 \leq t \leq 1$  is extremal for  $[\mu_t] \in T(S_0)$ . The arc  $t \rightarrow [\mu_t]$  is a geodesic from  $0$  to  $[\mu_1]$ , and  $d_T([\mu_t], 0) = td_T([\mu], 0)$ .

*Proof.* Since  $\mu$  represents a point in  $T(S_0)$ .

So  $\mu = (\mu \circ A) \frac{\bar{A}}{A}$  for every deck transformation  $A$ .

Therefore  $\mu_t$  also satisfies this condition, *i.e.*  $[\mu_t] \in T(S_0)$  is well-defined.

For  $g \in [\mu_t]$ , let  $g_0$  be the conformal map which is homotopic to  $f^{\mu_t} \circ g^{-1}$  and

$$h = f^\mu \circ (f^{\mu_t})^{-1} \circ g_0 \circ g.$$

Then  $h$  is homotopic to  $f^\mu$ , *i.e.*  $h \in [\mu_1]$ .

By theorem 3.6.1,  $K_1 \leq K_h \leq K_1^{1-t} K_g$ , *i.e.*  $K_g \geq K_1^t$ .

Therefore  $\mu_t$  is extremal for  $[\mu_t]$ .

And since  $d_T([\mu_t], [\mu_1]) \leq \frac{1}{2} \log K_{f^{\mu \circ (f^{\mu_t})^{-1}}} = (1-t)d_T([\mu], 0)$ .

So  $d_T(0, [\mu_t]) + d_T([\mu_t], [\mu_1]) = d_T(0, [\mu_1])$  for every  $0 \leq t \leq 1$ .

Hence  $t \rightarrow [\mu_t]$  is a geodesic. □

**Coro 4.3.1.**  $(T(S_0), d_T)$  is path connected.

*Proof.* For  $[\mu] \in T(S_0)$ , the arc  $t \rightarrow [\mu_t]$  is a path from 0 to  $[\mu_1]$ . □

**Thm 4.3.1.**  $(T(S_0), d_T)$  is complete.

*Proof.* Take a Cauchy sequence in  $(T(S_0), d_T)$  whose points are represented by Beltrami differential  $\mu_n$  and  $f_n \in [\mu_n]$ .

Fix a map  $f_i \in [\mu_i]$  such that for  $j \geq 1$ ,

$$\inf_{g_{i+j} \in [\mu_{i+j}]} \log K_{g_{i+j} \circ f_i^{-1}} < \frac{1}{2}.$$

Suppose  $f_i = f_1$  by passing to a subsequence and take  $f_n \in [\mu_n]$  such that

$$\log K_{f_n \circ f_1^{-1}} < \frac{1}{2}.$$

Choose  $f_k$  such that for  $j \geq 1$ ,

$$\inf_{g_{k+j} \in [\mu_{k+j}]} \log K_{g_{k+j} \circ f_k^{-1}} < \frac{1}{4}.$$

Suppose  $f_k = f_2$  by passing to a subsequence and take  $f_n \in [\mu_n]$  such that

$$\log K_{f_n \circ f_2^{-1}} < \frac{1}{4}.$$

Repeating this procedure gives a sequence  $\{f_n\}$  such that  $\{[f_n]\}$  is a subsequence of the Cauchy sequence and

$$\log K_{f_{n+1} \circ f_n^{-1}} < 2^{-n}.$$

So

$$\log K_{f_{n+j} \circ f_n^{-1}} \leq \sum_{i=1}^j 2^{-(n+i-1)} < 2^{-n+1}.$$

And the dilatation  $\mu_n$  of  $f_n$  satisfy that

$$\|\mu_{n+j} - \mu_n\|_\infty \leq 2 \left\| \frac{\mu_{n+j} - \mu_n}{1 - \bar{\mu}_n \mu_{n+j}} \right\|_\infty = 2 \tanh \left( \frac{1}{2} \log K_{f_{n+j} \circ f_n^{-1}} \right) < 2 \tanh 2^{-n}.$$

Therefore  $\{\mu_n\}$  is a Cauchy sequence in  $B^\infty(S_0)$ , *i.e.*  $\mu_n \rightarrow \mu$  as  $n \rightarrow \infty$  for some  $\mu \in B^\infty(S_0)$ . □

**Prop 4.3.4.** *Suppose  $g : S_0 \rightarrow X_0$  is quasiconformal, then  $T(S_0) \cong T(X_0)$ .*

*Proof.* Consider  $\phi : T(S_0) \rightarrow T(X_0), [S, f] \mapsto [S, f \circ g^{-1}]$ .

Then  $d([S_1, f_1], [S_2, f_2]) = d([S_1, f_1 \circ g^{-1}], [S_2, f_2 \circ g^{-1}])$ .  $\square$

**Def 4.3.5.**  $\text{Mod}_g = \{g : \Sigma_g \rightarrow \Sigma_g | g \text{ is diffeomorphism}\} / \sim$  and  $g_1 \sim g_2$  if  $g_1$  homotopic to  $g_2$ .

**Prop 4.3.5.**  $\text{Isom}(T_g) \cong \text{Mod}_g$ .

*Proof.* By proposition 4.3.4, every isometry  $T_g \rightarrow T_g$  corresponds to quasiconformal map  $\Sigma_g \rightarrow \Sigma_g$  and two maps  $f_1, f_2$  correspond to the same isometry iff  $[f_1] = [f_2] \in T_g$ , i.e.  $f_1 \simeq f_2$ .  $\square$

## 4.4 Douady-Earle extension

**Thm 4.4.1** (Schoen conjecture). *If  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  is quasi-symmetric, then there exists extension  $h : \mathbb{D} \rightarrow \mathbb{D}$  which is harmonic w.r.t. hyperbolic metric on  $\mathbb{D}$ .*

**Thm 4.4.2.** *If  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  is homeomorphism, then there exists a homeomorphic extension  $\phi_f : \mathbb{D} \rightarrow \mathbb{D}$  such that*

(1) (Conformally natural)  $A \circ \phi_f \circ B = \phi_{A \circ f \circ B}$  for  $A, B \in \text{Aut}(\mathbb{D})$

(2) If  $\varphi$  is quasi-symmetric, then  $\tilde{\varphi}$  is quasiconformal in  $\mathbb{D}$ .

Moreover, it is uniquely determined with the condition that

$$\int_{\mathbb{S}^1} f(\zeta) |d\zeta| = 0 \Leftrightarrow \phi_f(0) = 0$$

*Remark 4.4.1.* We take  $w = \phi_f(z)$  as the unique  $w \in \mathbb{D}$  such that

$$F(z, w) = \frac{1}{2\pi} \int_{\mathbb{S}^1} \left( \frac{f(\zeta) - w}{1 - \bar{w}f(\zeta)} \right) \left( \frac{1 - |z|^2}{|z - \zeta|^2} \right) |d\zeta| = 0.$$

It needs many technique of analysis to prove that the solution of  $F(z, w) = 0$  is unique for any  $z \in \mathbb{D}$  and such  $\phi_f$  satisfies the required statements, so we skip the detail.

**Lemma 4.4.1.** *Let  $S_0 = \mathbb{D}/\Gamma$ , consider the map*

$$\begin{aligned} \sigma : \text{Belt}(\mathbb{D}) &\longrightarrow \text{Belt}(\mathbb{D}) \\ \mu &\longmapsto \mu_{\phi_f \mu} \end{aligned}$$

where  $\varphi^\mu = f^\mu|_{\mathbb{S}^1}$ , then

(1)  $\sigma$  maps  $\text{Belt}(S_0)$  to itself

(2) there exists a continuous map  $s : T(S_0) \rightarrow \text{Belt}(S_0)$  such that  $s \circ \pi = \sigma$

(3)  $\pi \circ \sigma = \pi$ .

*Proof.* (1) Let  $B \circ f^\mu \circ A = f^\mu$  for some  $B \in \text{Aut}(\mathbb{D}), A \in \Gamma$ .

Then  $B \circ \varphi^\mu \circ A$ .

And since  $\phi$  is conformmally natural.

So  $B \circ \phi_{\varphi^\mu} \circ A = \phi_{\varphi^\mu}$ , i.e.  $\sigma(\mu) \in \text{Belt}(S_0)$ .

(2) Take  $s([\mu]) = \sigma(\mu)$ .

Suppose  $[\mu_1] = [\mu_2]$ , i.e.  $f^{\mu_1} \simeq f^{\mu_2}$ .

Then  $\varphi^{\mu_1} = \varphi^{\mu_2}$

So  $\phi_{\varphi^{\mu_1}} = \phi_{\varphi^{\mu_2}}$ , i.e.  $\sigma(\mu_1) = \sigma(\mu_2)$ .

Hence  $s$  is well-defined.

(3) Since  $\phi_{\varphi^\mu}|_{\mathbb{S}^1} = \varphi^\mu$ .

So  $\phi_{\varphi^\mu} \simeq f^\mu$ .

Hence  $\pi \circ \sigma(\mu) = [\mu]$ . □

**Thm 4.4.3.** *Let  $S_0 = \mathbb{D}/\Gamma$ , then  $T(S_0)$  is contractible.*

*Proof.* By the above lemma, since  $\pi \circ s \circ \pi = \pi \circ \sigma = \pi$ .

So  $\pi \circ s = \text{Id}$ .

Take  $H : T(S_0) \times [0, 1] \rightarrow T(S_0)$ ,  $([\mu], t) = [(1-t)s([\mu])]$ .

Then  $H(-, 0) = \text{Id}$  and  $H(-, 1) = 0$ .

Hence  $T(S_0)$  is contractible. □

## 4.5 Teichmüller space of torus

**Def 4.5.1.** Take  $T_\tau(z) = z + \tau$  and  $G_\tau = \langle T_1, T_\tau \rangle$ , then a torus is given by  $S_\tau = \mathbb{C}/G_\tau$ .

**Prop 4.5.1.**  $S_\tau = S_{\tau'}$  iff  $\tau' = A(\tau)$  for some  $A \in \text{PSL}(2, \mathbb{Z})$ .

**Lemma 4.5.1.** *Let  $\theta : G_\tau \rightarrow G_{\tau'}$  be an isomorphism, then there exists a homeomorphism  $f : S_\tau \rightarrow S_{\tau'}$  inducing  $\theta$ .*

*Proof.* Let  $T_{\omega_1} = \theta(T_1), T_{\omega_2} = \theta(T_\tau)$ .

Then consider the affine map  $f$  fixing 0 and mapping  $1, \tau$  to  $\omega_1, \omega_2$  resp.

Since  $\frac{\omega_2}{\omega_1} = A(\tau')$  for some  $A \in \text{PSL}(2, \mathbb{Z})$ .

So  $f$  projects to the map we desired. □

**Lemma 4.5.2.** *Let  $\tau, \tau' \in \mathbb{H}$ ,  $S_\tau$  and  $S_{\tau'}$  are conformally equivalent iff  $\tau' = A(\tau)$  for some  $A \in \text{SL}(2, \mathbb{Z})$ .*

*Proof.* Suppose  $S_\tau$  and  $S_{\tau'}$  are conformally equivalent and the conformal map lifts to  $f : \mathbb{C} \rightarrow \mathbb{C}$ .

Then  $f(0) = 0$ , i.e.  $f(z) = \alpha z$ .

So  $(\alpha, \alpha\tau)$  is a base of  $S_{\tau'}$ , i.e.  $\tau' = A(\tau)$  for some  $A \in \text{SL}(2, \mathbb{Z})$ .

Conversely, by proposition 4.5.1,  $S_\tau = S_{\tau'}$ . □

**Thm 4.5.1.** *In each homotopy class of sense-preserving homeomorphisms between tori, the extremal map can be lifted to an affine map.*

*Proof.* Suppose  $f : S_\tau \rightarrow S_{\tau'}$  and the lift  $\tilde{f} : \mathbb{C} \rightarrow \mathbb{C}$  satisfies that  $f(0) = 0$  and

$$\tilde{f}(z + m + n\tau) = \tilde{f}(z) + m\omega_1 + n\omega_2,$$

we denote the set of such  $\tilde{f}$  by  $F$ .

Then there exists a unique affine map

$$g(z) = \frac{\omega_2 - \omega_1\bar{\tau}}{\tau - \bar{\tau}}z + \frac{\omega_1\tau - \omega_2}{\tau - \bar{\tau}}\bar{z}$$

in  $F$ .

Suppose  $f$  is  $K$ -quasiconformal, let  $f_k(z) = \frac{\tilde{f}(kz)}{k}$

Then  $f_k$  converges to an affine map in  $F$ , i.e.  $f_k \rightarrow g$ .

So  $K_g \leq K$ .

Now suppose that  $K = K_g$ , assume  $g(x + iy) = Kx + iy$  WLOG.

Since  $\tilde{f}, g \in F$ , there exists  $M$  such that

$$|\tilde{f}(z) - g(z)| \leq M.$$

So we obtain

$$\int_0^r |\tilde{f}_x(x + iy)| dx \geq \left| \int_0^r \tilde{f}_x(x + iy) dx \right| \geq Kr - 2L$$

Consider  $R = [0, r] \times [0, r]$ , then

$$\int_R |\tilde{f}_x| dx dy \geq Kr^2 - 2Lr.$$

As  $r \rightarrow \infty$ , the number of period parallelograms  $P$  meeting  $R$  is about  $\frac{r^2(1+o(1))}{A(P)}$ , and so

$$\int_P |\tilde{f}_x| dx dy \geq \lim_{r \rightarrow \infty} \frac{r^2(K + o(1))}{r^2(1 + o(1))} A(P) = KA(P).$$

On the other hand,  $|f_x|^2 \leq KJ$ , therefore

$$(KA(P))^2 \leq A(P) \int_P |\tilde{f}_x|^2 dx dy \leq KA(P)A(\tilde{f}(P)).$$

And since  $\tilde{f}(P)$  is the fundamental domain of  $S_{\tau'}$ , i.e.  $A(\tilde{f}(P)) = A(g(P)) = KA(P)$ .

Hence  $|\tilde{f}_x|^2 = KJ$  almost everywhere, i.e.  $\tilde{f} = g$  is affine.  $\square$

**Lemma 4.5.3.** *Let  $\theta : G_\tau \rightarrow G_{\tau'}$  be an isomorphism generated by  $K$ -quasiconformal map  $f$ , then*

$$d_{\mathbb{H}} \left( \tau, \frac{\tilde{f}(\tau)}{\tilde{f}(1)} \right) \leq \frac{1}{2} \log K$$

*the equality holds iff  $\tilde{f}$  is affine.*

*Proof.* WLOG, we assume  $\frac{f(\tau)}{f(1)} = \tau'$ .

Let  $g$  homotopic to  $f$  and  $\tilde{g} = \lambda(z + \mu\bar{z})$  is affine, then

$$\mu = \frac{\tau - \tau'}{\tau' - \bar{\tau}}.$$

So we obtain

$$\log K_g = \log \frac{1 + |\mu|}{1 - |\mu|} = \tanh \left( \frac{|\tau - \tau'|}{|\tau' - \bar{\tau}|} \right) = 2d_{\mathbb{H}}(\tau, \tau').$$

And by the above theorem,  $K_g \geq K_f$ .

Hence we complete the proof.  $\square$

**Thm 4.5.2.** *Teichmüller space of torus is isometric to  $\mathbb{H}$  with hyperbolic metric.*

*Proof.* Consider  $\psi : T(S_\tau) \rightarrow \mathbb{H}$  given by

$$\psi([f]) = \frac{\tilde{f}(\tau)}{\tilde{f}(1)}.$$

By three lemmas above, we can easily check that  $\psi$  is bijective.

Consider  $[f_1], [f_2] \in T(S_\tau)$ , let  $f : f_1(S_\tau) \rightarrow f_2(S_\tau)$  be the extremal homotopic to  $f_2 \circ f_1^{-1}$ . Then by lemma 4.5.3,

$$d_T([f_1], [f_2]) = \frac{1}{2} \log K_f = d_{\mathbb{H}} \left( \frac{\tilde{f}_1(\tau)}{\tilde{f}_1(1)}, \frac{\tilde{f}_2(\tau)}{\tilde{f}_2(1)} \right) = d_{\mathbb{H}}(\psi([f_1], [f_2])).$$

□

## 4.6 Teichmüller theorem

**Def 4.6.1.** Let  $S$  be a Riemann surface and  $\nu \in \text{Belt}(S)$ , we say  $\nu$  annihilates  $QD^1(S)$  if

$$\int_S \gamma \phi = 0$$

for all  $\phi \in QD^1(S)$ .

**Def 4.6.2.** Consider the linear functional

$$\Lambda_\mu(\phi) = \int_M \mu(z) \phi(z) dx dy.$$

We say  $\mu$  and  $\nu$  are infinitesimally equivalent, or  $\mu \sim_* \nu$  if  $\Lambda_\mu(\phi) = \Lambda_\nu(\phi)$  for  $\phi \in QD^1(S)$ . The infinitesimally equivalent class of  $\mu$  is denoted by  $[\mu]_*$ .

An element  $\mu$  is said to be infinitesimally extremal if for all  $\nu \in [\mu]_*$ ,  $\|\mu\|_\infty \leq \|\nu\|_\infty$ .

**Thm 4.6.1** (Hamilton–Krushkal). *Suppose  $\mu \in \text{Belt}(S)$  is extremal in  $[\mu] \in T(S)$ , then  $\mu$  is infinitesimally extremal.*

*Proof.* Let  $\|\mu\|_\infty = k$ ,  $\alpha \in [\mu]_*$  and  $\|\alpha\|_\infty = k_1$ .

Suppose  $k_1 < k$ , let  $\nu = \mu - \alpha$

Then  $\nu$  annihilates  $QD^1(S)$ .

By theorem 4.1.2, take  $[\delta(t)] = [t\nu] \in T(S)$  such that  $\|\delta(t)\|_\infty \leq 12t^2$ , let  $\nu_t = t\nu - \delta(t)$ .

So  $[\nu_t] = [0]$  in  $T(S)$ .

Moreover, take  $f^\lambda = f^\mu \circ (f^{\nu_t})^{-1}$ , then

$$\begin{aligned} |\lambda \circ f^{\nu_t}| &= \left| \frac{\mu - \nu_t}{1 - \mu \bar{\nu}_t} \right| \\ &= \sqrt{\frac{|\mu|^2 - 2 \operatorname{Re}(\mu \bar{\nu}_t) + |\nu_t|^2}{1 - 2 \operatorname{Re}(\mu \bar{\nu}_t) + |\nu_t \mu|^2}} \\ &= \sqrt{|\mu|^2 - 2(1 - |\mu|^2) \operatorname{Re}(\mu \bar{\nu}) + O(t^2)} \\ &= |\mu| - \frac{1 - |\mu|^2}{|\mu|} \operatorname{Re}(\mu \bar{\nu}) t + O(t^2) \end{aligned}$$

Consider  $E_1 = \left\{ z \in S \mid |\mu(z)| \leq \frac{k+k_1}{2} \right\}$ ,  $E_2 = S \setminus E_1$ .

On  $E_1$ ,  $|\lambda(f^{\nu_t}(z))| \leq \frac{k+k_1}{2} + O(t) < k - C_1 t$  for small  $t > 0$ .

On  $E_2$ , since

$$\operatorname{Re}(\mu\bar{v}) = t \operatorname{Re}(|\mu|^2 - \mu\bar{\alpha}) \geq t|\mu| \operatorname{Re}(|\mu| - |\alpha|).$$

So we deduce

$$\frac{1 - |\mu|^2}{|\mu|} \operatorname{Re}(\mu\bar{v}) \geq (1 - |\mu|^2)(|\mu| - |\alpha|) \geq \frac{1}{2}(1 - k^2)(k - k_1)$$

Therefore  $|\lambda(z)| < k - C_2t$  for small  $t > 0$  and  $z \in E_2$ .

Hence  $|\lambda(z)| < k - Ct$  for some constant  $C$  and  $z \in S$ , contradiction!  $\square$

**Thm 4.6.2** (Reich-Strebel inequality). *Suppose  $f^\mu : S \rightarrow S$  and  $[\mu] = 0$ , then for  $\phi \in \operatorname{QD}^1(S)$ ,*

$$\|\phi\|_1 \leq \int_S |\phi| \frac{|1 + \mu \frac{\phi}{|\phi|}|^2}{1 - |\mu|^2} dx dy$$

*Remark 4.6.1.* The proof uses the theory of trajectories of quadratic differentials, so we omit the proof.

**Exam 4.6.1.** *Consider  $\lambda = k \frac{|\phi|}{\phi}$  and  $\phi_0 = -\phi$ , then*

$$\int_S |\phi| \frac{|1 - \mu \frac{\phi}{|\phi|}|^2}{1 - |\mu|^2} dx dy = \int_S |\phi| \frac{1 - k}{1 + k} dx dy.$$

*So the Reich-Strebel inequality cannot hold, i.e.  $[\lambda] \neq 0$ .*

We now proof the special case of Reich-Strebel inequality for square torus.

**Prop 4.6.1** (Grötzsch). *Let  $S$  be a square torus,  $f^\mu : S \rightarrow S$  and  $[\mu] = 0$ , then for  $\phi \in \operatorname{QD}^1(S)$ ,*

$$\|\phi\|_1 \leq \int_S |\phi| \frac{|1 + \mu \frac{\phi}{|\phi|}|^2}{1 - |\mu|^2} dx dy$$

*Proof.* Take  $\gamma_y(t) = (t, y)$ .

Then  $l(f(\gamma_y)) \geq l(\gamma_y) = 1$ , so

$$\begin{aligned} \int_0^1 1 dy &\leq \int_0^1 \int_0^1 |f_z(x, y)| |1 + \mu(x, y)| dx dy \\ &= \int_S |f_z| |1 + \mu| dx dy \\ &= \int_S \frac{|f_z| |1 + \mu|}{\sqrt{|f_z|^2 - |f_{\bar{z}}|^2}} \sqrt{|f_z|^2 - |f_{\bar{z}}|^2} dx dy \\ &\leq \int_S \frac{|1 + \mu|^2}{1 - |\mu|^2} dx dy \cdot \int_S (|f_z|^2 - |f_{\bar{z}}|^2) dx dy \\ &= \int_S \frac{|1 + \mu|^2}{1 - |\mu|^2} dx dy \end{aligned}$$

And since  $\operatorname{QD}^1(S) = \mathbb{C}dz^2$ .

So we conclude the proof.  $\square$

**Coro 4.6.1** (main inequality). *Suppose  $f : S \rightarrow f(S)$  and  $\tilde{f} : f(S) \rightarrow S$  have Beltrami differentials  $\mu$  and  $\mu_1$  resp. and  $\tilde{f} \circ f$  is homotopic to the identity, then for any  $\phi \in \text{QD}^1(S)$*

$$\|\phi\|_1 \leq \int_S \frac{\left|1 + \mu \frac{\phi}{|\phi|}\right|^2}{1 - |\mu|^2} \frac{\left|1 + \theta \mu_1(f) \frac{\phi}{|\phi|}\right|^2}{1 - |\mu_1(f)|^2} |\phi| dx dy.$$

Here

$$\theta = p \left(1 + \frac{\overline{\mu\phi}}{|\phi|}\right) \left(1 + \frac{\mu\phi}{|\phi|}\right)^{-1}, p = \frac{\overline{f_z}}{f_z}$$

*Proof.* The Beltrami differential  $\lambda$  of  $\tilde{f} \circ f$  is

$$\lambda = \frac{\mu + \mu_1(f)p}{1 + \overline{\mu}\mu_1(f)p}$$

So by Reich-Strebel inequality,

$$\begin{aligned} \|\phi\|_1 &\leq \int_S |\phi| \frac{\left|1 + \lambda \frac{\phi}{|\phi|}\right|^2}{1 - |\lambda|^2} dx dy \\ &= \int_S |\phi| \frac{\left|1 + \overline{\mu}\mu_1(f)p + (\mu + \mu_1(f)p) \frac{\phi}{|\phi|}\right|^2}{\left|1 + \overline{\mu}\mu_1(f)p\right|^2 - \left|\mu + \mu_1(f)p\right|^2} dx dy \\ &= \int_S |\phi| \frac{\left|1 + \mu \frac{\phi}{|\phi|}\right|^2 \left|1 + \frac{\overline{\mu} + \frac{\phi}{|\phi|}}{1 + \mu \frac{\phi}{|\phi|}} \mu_1(f)p\right|^2}{(1 - |\mu|^2) (1 - |\mu_1(f)p|^2)} dx dy \\ &= \int_S \frac{\left|1 + \mu \frac{\phi}{|\phi|}\right|^2}{1 - |\mu|^2} \frac{\left|1 + \theta \mu_1(f) \frac{\phi}{|\phi|}\right|^2}{1 - |\mu_1(f)|^2} |\phi| dx dy \end{aligned}$$

□

**Thm 4.6.3.** *Suppose  $f : S_0 \rightarrow S$  with Beltrami differential of Teichmüller type, i.e.  $\mu = k \frac{\overline{\phi}}{|\phi|}$  for some  $\phi \in \text{QD}^1(S)$ , and  $g : S_0 \rightarrow S$  is quasiconformal such that  $g^{-1} \circ f$  homotopic to the identity, then either there exists a set of positive measure on  $S_0$  for which  $|\mu_g(z)| > k$  or  $\mu_g(f) = \mu$  almost everywhere.*

*Proof.* Let  $\nu(z) = \text{Belt}(g^{-1})(f(z))$  and WLOG, we assume  $\|\phi\|_1 = 1$ .

If  $\|\mu_g\|_\infty = \|\nu\|_\infty \leq k$ , then

$$\begin{aligned} 1 &\leq \int_S \frac{\left|1 - \mu \frac{\phi}{|\phi|}\right|^2}{1 - |\mu|^2} \frac{\left|1 - \theta \nu \frac{\phi}{|\phi|}\right|^2}{1 - |\nu|^2} |\phi| dx dy \\ &= \int_S \frac{1 - k}{1 + k} \frac{\left|1 - \frac{\overline{f_z} \nu}{f_z} \frac{\phi}{|\phi|}\right|^2}{1 - |\nu|^2} |\phi| dx dy \\ &\leq \frac{1}{K} \int_S \frac{1 + |\nu|^2}{1 - |\nu|^2} |\phi| dx dy \\ &\leq \frac{1}{K} \frac{1 + \|\nu\|_\infty}{1 - \|\nu\|_\infty} \|\phi\|_1 \leq 1 \end{aligned}$$

So each inequality must be equality, i.e.

$$\left|1 + \mu_g(f) \frac{\phi}{|\phi|}\right|^2 = \left|1 - \frac{\overline{f_z} \nu}{f_z} \frac{\phi}{|\phi|}\right|^2 = 1 + |\nu|^2, |\nu| \equiv k$$

hold almost everywhere.

Hence  $\mu_g(f) = \mu$  almost everywhere.  $\square$

**Thm 4.6.4** (Teichmüller existence theorem). *Let  $X$  Riemann surface with  $\dim \text{QD}^1(X) < +\infty$ , then every point in  $T(X)$  has a uniquely extremal representative, which is of Teichmüller type.*

*Proof.* Take  $[\mu] \in T(X)$  such that  $\mu$  is extremal in  $[\mu]$ , consider the linear functional

$$\lambda(\phi)(\phi) = \int_X \mu\phi.$$

Then there exists  $\phi_0 \in \text{QD}^1(X)$  such that  $\|\lambda\| = \lambda(\phi_0)$  and  $\|\phi_0\| = 1$ .

By Hamilton-Krushkal condition,  $\mu$  is infinitesimal extremal, *i.e.*

$$\|\mu\|_\infty = \|\lambda\| = \int_X \mu\phi.$$

So  $\mu = k \frac{\bar{\phi}_0}{|\phi_0|}$ .

By theorem 4.6.3,  $\mu$  is the unique extremal.  $\square$

**Prop 4.6.2.** *Let  $\tau_1, \tau_2 \in T(X)$ , then there exists a geodesic between  $\tau_1, \tau_2$  and moreover, if  $\dim \text{QD}^1(X) < +\infty$ , then this geodesic is unique.*

*Proof.* By proposition 4.3.3, the geodesic exists.

And if  $\dim \text{QD}^1(X) < +\infty$ , by the Teichmüller existence theorem, the geodesic is unique.  $\square$

*Remark 4.6.2.* For  $\mu = k \frac{\bar{\phi}}{|\phi|}$ , the unique geodesic from 0 to  $\mu$  is given by

$$t \mapsto \left[ \tanh(t) \frac{\bar{\phi}}{|\phi|} \right].$$

So the tangent bundle of  $T(X)$  is given by  $\text{Belt}(X)/\sim_*$ , which is the dual space of  $\text{QD}^1(X)$ . Moreover,  $T(X)$  is has a Finsler norm

$$\|[\mu]_*\|_{\text{Fin}} = \inf_{\nu \in [\mu]_*} \|\nu\|_\infty.$$

**Def 4.6.3.** Consider  $\mu \in \text{Belt}(\mathbb{H}/\Gamma)$ , we can extend  $\mu$  on  $\mathbb{C}$  with  $\mu(z) = \overline{\mu(\bar{z})}$ , then  $f^\mu(z) = \overline{f^\mu(\bar{z})}$ , *i.e.* we can induce  $f^\mu : X \rightarrow X^1 \text{mu} = \mathbb{H}/\Gamma^\mu$ .

On the other hand, if we extend  $\mu$  on  $\mathbb{C}$  such that  $\mu(z) = 0$  for  $\text{Im } z < 0$ , we can define  $f_\mu : \mathbb{C} \rightarrow \mathbb{C}$ .

## 4.7 Schwartzian derivative

**Def 4.7.1.** If  $f$  is a conformal map, or more generally, for holomorphic map with  $f'(z) \neq 0$ , we define

$$(Sf)(z) = 6 \lim_{w \rightarrow z} \frac{\partial^2}{\partial z \partial w} \log \frac{f(z) - f(w)}{z - w}.$$

*Remark 4.7.1.* Notice that

$$\int_{z_1}^{z_2} \int_{w_1}^{w_2} \frac{\partial^2}{\partial z \partial w} \log \frac{f(z) - f(w)}{z - w} dw dz = \log \frac{\text{Cr}(f(z_1), f(w_2), f(z_2), f(w_1))}{\text{Cr}(z_1, w_2, z_2, w_1)}$$

So In some sence,  $S(f)$  measures the amounts by which  $f$  distorts the cross ratios.

**Prop 4.7.1.**

$$(Sf)(z) = \frac{f'''}{f'}(z) - \frac{3}{2} \left( \frac{f''}{f'} \right)^2 (z).$$

*Proof.*

$$\begin{aligned} (Sf)(z) &= 6 \lim_{t \rightarrow 0} \frac{f'(z)f'(z+t)}{(f(z) - f(z+t))^2} - \frac{1}{t^2} \\ &= 6 \lim_{t \rightarrow 0} \frac{f'(z) \left( f'(z) + f''(z)t + \frac{1}{2}f'''(z)t^2 + o(t^2) \right)}{t^2 \left( f'(z) + \frac{1}{2}f''(z)t + \frac{1}{6}f'''(z)t^2 + o(t^2) \right)^2} - \frac{1}{t^2} \\ &= 6 \lim_{t \rightarrow 0} \frac{f'(z)}{t^2} \cdot \frac{f'(z) + f''(z)t + \frac{1}{2}f'''(z)t^2 + o(t^2)}{f'(z)^2 + f'(z)f''(z)t + \left( \frac{1}{4}f''(z)^2 + \frac{1}{3}f'(z)f'''(z) \right) t^2 + o(t^2)} - \frac{1}{t^2} \\ &= 6 \lim_{t \rightarrow 0} \frac{1}{t^2} \left( 1 + \frac{f''}{f'}(z)t + \frac{1}{2} \frac{f'''}{f'}(z)t^2 + o(t^2) \right) \\ &\quad \cdot \left( 1 - \frac{f''}{f'}(z)t + \left( \left( \frac{f''}{f'} \right)^2 (z) - \frac{1}{4} \left( \frac{f''}{f'} \right)^2 (z) - \frac{1}{3} \frac{f'''}{f'}(z) \right) t^2 + o(t) \right) - \frac{1}{t^2} \\ &= 6 \lim_{t \rightarrow 0} \frac{1}{t^2} \left( 1 + \left( \frac{1}{6} \frac{f'''}{f'}(z) - \frac{1}{4} \left( \frac{f''}{f'} \right)^2 (z) \right) t^2 + o(t) \right) - \frac{1}{t^2} \\ &= \frac{f'''}{f'}(z) - \frac{3}{2} \left( \frac{f''}{f'} \right)^2 (z) \end{aligned}$$

□

**Prop 4.7.2.**

$$S(f \circ g) = ((Sf) \circ g) \cdot (g')^2 + Sg$$

and  $Sf \equiv 0$  iff  $f$  is a Möbius transformation. In particular, for Möbius transformation  $A$ ,

$$S_{f \circ A} = (Sf \circ A) \cdot (A')^2.$$

*Proof.*

$$\begin{aligned} S(f \circ g)(z) &= 6 \lim_{w \rightarrow z} \frac{(f \circ g)'(z)(f \circ g)'(w)}{(f(g(z)) - f(g(w)))^2} - \frac{1}{(z-w)^2} \\ &= 6 \lim_{w \rightarrow z} \frac{f'(g(z))f'(g(w))}{(f(g(z)) - f(g(w)))^2} g'(z)g'(w) - \frac{1}{(z-w)^2} \\ &= (Sf)(g(z)) \cdot g'(z)^2 + 6 \lim_{w \rightarrow z} \frac{g'(z)g'(w)}{(g(z) - g(w))^2} - \frac{1}{z-w} \\ &= (Sf)(g(z)) \cdot g'(z)^2 + (Sg)(z) \end{aligned}$$

Moreover,  $Sf \equiv 0$  iff  $f$  preserved cross-ratio, i.e.  $f$  is a Möbius transformation. □

**Def 4.7.2.** Define

$$\begin{aligned} \tilde{B}: \text{Belt}(\mathbb{H}/\Gamma) &\longrightarrow \text{QD}(\mathbb{H}^-/\Gamma) \\ \mu &\longmapsto S(f_\mu|_{\mathbb{H}^-}) \end{aligned}$$

Then the Bers embedding is given by

$$\begin{aligned} B: \text{Belt}(\mathbb{H}/\Gamma)/\sim &\longrightarrow \text{QD}(\mathbb{H}^-/\Gamma) \\ [\mu] &\longmapsto \tilde{B}(\mu) \end{aligned}$$

*Proof.* Take  $\mu \in \text{Belt}(\mathbb{H}/\Gamma)$ .

Then  $f_\mu \circ \gamma \circ f_\mu^{-1}$  is a Möbius transformation for any  $\gamma \in \Gamma$ , so

$$S(f_\mu|_{\mathbb{H}^-}) = S(f_\mu \circ \gamma \circ f_\mu^{-1} \circ f_\mu|_{\mathbb{H}^-}) = S(f_\mu \circ g|_{\mathbb{H}^-}) = (S(f_\mu|_{\mathbb{H}^-}) \circ \gamma) \cdot (\gamma')^2$$

Hence  $S(f_\mu|_{\mathbb{H}^-}) \in \text{QD}(\mathbb{H}^-/\Gamma)$  is well-defined.

Moreover, by proposition 4.3.1, if  $\mu$  homotopic to  $\nu$ , then  $f_\mu|_{\mathbb{H}^-} = f_\nu|_{\mathbb{H}^-}$ .  $\square$

**Prop 4.7.3** (Nehari). *Let  $\rho$  be the hyperbolic metric on  $X = \mathbb{H}/\Gamma$ , for every  $\mu \in \text{Belt}(\mathbb{H}/\Gamma)$ ,*

$$\|\rho^{-2}S(f_\mu)\|_\infty \leq \frac{3}{2}.$$

*Proof.*  $\rho^{-2}S(f_\mu)$  is invariant under Möbius transformations.

WLOG, we can only estimate  $\rho(0)^{-2}S(f_\mu)(0)$  and assume  $f'(0) = 1$ .

Consider the function

$$\frac{1}{f(z^{-1})} = z + \sum_{n=0}^{\infty} b_n z^{-n}.$$

By the Gronwall area theorem, we obtain that

$$|b_1| = |a_2^2 - a_3| \leq 1.$$

So  $|S_f(0)| = 6|a_3 - a_2^2| \leq 6$ .

Hence we complete the proof.  $\square$

**Exam 4.7.1.** *Take  $f(z) = z^2$ , then  $Sf = -\frac{3}{2z^2}$ , so the bound can be attained.*

# Chapter 5

## Additional topics

This chapter are some additional topics about Teichmüller theory, most of the proposition has no proof. You can try to find them if you are interested.

### 5.1 Bicanonical embedding

**Def 5.1.1.** Suppose  $S$  is a Riemann surface and  $a_1, \dots, a_n \in S$ , a divisor is

$$D = \sum_{i=1}^n m_i a_i,$$

and its degree is given by

$$\deg(D) = \sum_{i=1}^n m_i.$$

**Def 5.1.2.** A holomorphic map  $f : S \rightarrow \mathbb{C}P^1$  is called meromorphism and we define

$$\text{ord}_a(f) = \begin{cases} k & f \text{ has a zero of order } k \text{ at } a \\ -k & f \text{ has a pole of order } k \text{ at } a \end{cases}$$

The principal divisor of  $f$  is given by

$$(f) = \sum_{a \in S} \text{ord}_a(f) a$$

**Prop 5.1.1.**  $\deg((f)) = 0$ .

**Def 5.1.3.** Let  $L(D)$  be the set of meromorphic functions on  $S$  such that  $(f) + D \geq 0$  and we denote  $l(D) = \dim L(D)$ .

**Def 5.1.4.** If  $\omega$  is abelian differential, *i.e.* meromorphic 1-form, we say  $(\omega)$  is canonical.

If  $\phi \in \text{QD}(s)$ , we say  $(\phi)$  is bicanonical.

**Prop 5.1.2.**  $\deg(\omega) = -\chi(S)$ .

**Def 5.1.5.**  $I(D)$  is the space of Abelian differentials with  $(\omega) \geq D$  and  $i(D) = \dim I(D)$ .

**Thm 5.1.1** (Riemann-Roch).

$$l(D) - i(D) = \deg(D) + 1 - g.$$

**Thm 5.1.2.**  $S$  is a closed Riemann surface with genus  $g \geq 2$ , then  $\dim \text{QD}(S) = 3g - 3$ .

*Proof.* Let  $\omega$  be an Abelian differential and  $D = 2(\omega)$ , then

$$\text{QD}(S) \cong \{\text{meromorphic } f : S \rightarrow \mathbb{C}\mathbb{P}^1 \mid (f) + D \geq 0\} = L(D)$$

So by Riemann-Roch,

$$\dim \text{QD}(S) = l(D) = i(D) + \deg(D) + 1 - g = i(D) + 3g - 3.$$

And since for any Abelian differential  $\alpha$ ,  $\deg(\alpha) = \deg(\omega)$ , *i.e.* we cannot have  $(\alpha) \geq 2(\omega)$ . Hence  $i(D) = 0$ , *i.e.*  $\dim \text{QD}(S) = 3g - 3$ .  $\square$

**Prop 5.1.3** (canonical embedding). *Let  $\omega_1, \dots, \omega_g$  be the basis of  $\Omega(S)$ , i.e. the set of all abelian differentials, if  $S$  is non-hyperelliptic, then*

$$f : S \rightarrow \mathbb{C}\mathbb{P}^{g-1}, f = [\omega_1 : \dots : \omega_g]$$

*is an embedding.*

**Prop 5.1.4** (bicanonical embedding). *Let  $\phi_1, \dots, \phi_{3g-3}$  be the basis of quadratic differential forms on  $S$ , if  $g > 2$ , then*

$$f : S \rightarrow \mathbb{C}\mathbb{P}^{3g-4}, f = [\phi_1 : \dots : \phi_{3g-3}]$$

*is an embedding.*

## 5.2 Automorphism of $T_g$

In this section, we assume  $g > 2$ .

For a biholomorphism (isometry)  $I : T_g \rightarrow T_g$  with  $\tau \in T_g$ , there is an induced isometric

$$\begin{aligned} I^* : T_{I(\tau)}^* T_g &\rightarrow T_\tau^* T_g \\ L : \text{QD}(Y) &\rightarrow \text{QD}(X) \end{aligned}$$

where  $X \in \tau, Y \in I(\tau)$ .

**Thm 5.2.1.** *Suppose  $(X, \mu), (Y, \nu)$  are two finite measure spaces,  $f_1, \dots, f_k \in L^p(\mu)$  and  $g_1, \dots, g_k \in L^p(\nu)$  such that*

$$\int_X \left| 1 + \sum \lambda_j f_j \right|^p d\mu = \int_Y \left| 1 + \sum \lambda_j g_j \right|^p d\nu$$

*for any  $\lambda = (\lambda_1, \dots, \lambda_k) \in \mathbb{C}^k$ , then*

$$\langle 1, f_1, \dots, f_k \rangle \leftrightarrow \langle 1, g_1, \dots, g_k \rangle$$

*is isometric w.r.t.  $L^p$ -norm.*

*Moreover, if  $0 < p < \infty$  and  $p$  is not even integer, then  $\mu(F^{-1}(E)) = \nu(G^{-1}(E))$  for every Borel set  $E \subset \mathbb{C}^k$  with  $F = (f_1, \dots, f_k), G = (g_1, \dots, g_k)$ .*

**Prop 5.2.1.** *If  $L : \text{QD}(X) \rightarrow \text{QD}(Y)$  is a complex linear isomorphism w.r.t.  $L^1$ -norm, then there exists biholomorphic  $f : Y \rightarrow X$  such that  $L(\phi) = f^* \phi$ .*

*Proof.* Let  $\phi_0, \dots, \phi_k$  be the basis of  $\text{QD}^1(X)$  and  $\psi_0, \dots, \psi_k$  be the basis of  $\text{QD}^1(Y)$ .

Then  $L(\phi_i) = \psi_i$ .

Consider  $f_i = \frac{\phi_i}{\phi_0}, g_i = \frac{\psi_i}{\psi_0}$ , so

$$\int_X \left| 1 + \sum \lambda_i f_i \right| d\mu = \int_Y \left| 1 + \sum \lambda_i g_i \right| d\nu$$

where  $d\mu = |\phi_0| dx dy, d\nu = |\psi_0| dx dy$ .

By the above theorem, this is possible only when there exists biholomorphic  $f : Y \rightarrow X$  such that  $f^* \mu = \nu$ .  $\square$

### 5.3 Quasi-Fuchsian group

Consider Fuchsian group  $\Gamma$  and  $\mathbb{H}^+/\Gamma = X, \mathbb{H}^-/\Gamma = \bar{X}$  in  $\mathbb{S}^2$ .  
Take  $\mu^+ \in \text{Belt}(X), \mu^- \in \text{Belt}(\bar{X})$  and

$$\mu = \begin{cases} \mu^+(z) & z \in \mathbb{H}^+ \\ \mu^-(z) & z \in \mathbb{H}^- \end{cases}$$

Then we obtain a homeomorphism  $f^\mu : \mathbb{S}^2 \rightarrow \mathbb{S}^2$ .

**Prop 5.3.1.**  $\mu$  is equivariant under  $\Gamma$ , i.e.  $(\mu \circ A) \frac{\bar{A}}{A} = \mu$  on  $\mathbb{S}^2$ .

So there exists a group of Möbius transformation  $\Gamma^\mu$  such that  $f^\mu$  conjugates  $\Gamma$  to  $\Gamma^\mu$ .

**Def 5.3.1.**  $\mathbb{H}^3/\Gamma$  is called hyperbolic Fuchsian 3-manifold.

$\Gamma^\mu$  is called quasi-Fuchsian group and  $\mathbb{H}^3/\Gamma^\mu$  is called hyperbolic quasi-Fuchsian 3-manifold.

*Remark 5.3.1.* We say  $\Gamma^\mu$  is “quasi-Fuchsian” because  $f^\mu(\mathbb{R})$  is a quasicircle and  $\Gamma^\mu$  preserve this curve since  $\Gamma^\mu = f^\mu \Gamma (f^\mu)^{-1}$ .

Notice that if  $\mu^+ \sim \nu^+, \mu^- \sim \nu^-$ , then  $\mu \sim \nu$  and  $\Gamma^\mu \cong \Gamma^\nu$ . So we have a map

$$\begin{aligned} \mathcal{B} : T(X) \times T(\bar{X}) &\longrightarrow \{\text{Quasi-Fuchsian 3-manifolds}\} \\ ([\mu^+], [\mu^-]) &\longmapsto \mathbb{H}^3/\Gamma^\mu \end{aligned}$$

**Prop 5.3.2.** Fuchsian and quasi-Fuchsian 3-manifolds are homeomorphic to  $\Sigma_g \times (0, 1)$  for some  $g$ .

*Proof.* Consider Fuchsian group  $\Gamma$  and denote  $H = \mathbb{D}^3 \cap \mathbb{R}^2$ .

Then on every equi-distance surface of  $H$ , the quotient is homeomorphic to  $\mathbb{H}^2/\Gamma$ .

Hence  $\mathbb{H}^3/\Gamma \cong (\mathbb{H}^2/\Gamma) \times \mathbb{R} = \Sigma_g \times (0, 1)$ . □

**Def 5.3.2.** Suppose  $M$  is a hyperbolic 3-manifold, the convex core  $C(M)$  is the smallest closed convex subset of  $M$  containing every closed geodesic on  $M$ .

**Prop 5.3.3.** Suppose  $M = \mathbb{H}^3/\Gamma$ , then  $C(M)$  is the quotient of the convex hull of the limit set  $\Lambda(\Gamma) \subset \mathbb{S}^2$  by  $\Gamma$ .

*Proof.* Since every closed geodesic on  $M$  can be lifted to an axis  $\gamma$  of  $\Gamma$ .

So the endpoints of  $\gamma$  lie in  $\Lambda(\Gamma)$ , i.e.  $\gamma \subset \text{Ch}(\Lambda(\Gamma))$ .

Conversely, for  $\xi \in \Lambda(\Gamma)$ , there exists a sequence of  $g_n \in \Gamma$  whose fixed point approach  $\xi$ .

So the axis  $\gamma_n$  of  $g_n$  converges to a geodesic  $\gamma$  start at  $\xi$ .

Therefore the quotient of  $\gamma_n$  converges to the quotient  $\gamma$ , i.e. it is contained in  $C(M)$ .

Hence  $C(M) = \text{Ch}(\Lambda(\Gamma))/\Gamma$ . □

**Thm 5.3.1.** If  $M$  is a hyperbolic 3-manifold which is homeomorphic to  $\Sigma_g \times (0, 1)$  and  $C(M)$  is compact in  $M$ , then  $M$  is quasi-Fuchsian.

**Prop 5.3.4.**  $p \in \mathbb{S}^2 \setminus \Lambda(\Gamma)$  iff there exists a neighborhood  $U$  of  $p$  such that

$$\{A|_U : A \in \Gamma\}$$

is a normal family.

## 5.4 Complex dynamics

**Def 5.4.1.** Rational maps are of the form  $f : \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{C}\mathbb{P}^1, z \mapsto \frac{P(z)}{Q(z)}$  where  $P, Q$  are relatively prime polynomial, its degree is given by  $\deg f = \max\{\deg P, \deg Q\}$ .

Let  $\text{Rat}_d$  be the space of degree  $d$  rational maps.

Now take  $f \in \text{Rat}_d$ , the iterations of  $f$  is a sequence  $\{f^n\}_{n \in \mathbb{Z}}$ .

**Def 5.4.2.** The orbit of iteration is defined as

$$\begin{aligned} O^+(z) &= \{f^n(z) | n \in \mathbb{N}\}, \\ O^-(z) &= \{f^n(z) | n \in \mathbb{Z}^-\}, \\ O_{\text{grand}}(z) &= \bigcup_{n \geq 0} O^-(f^n(z)). \end{aligned}$$

**Def 5.4.3.** Two maps are called Möbius equivalent if  $g = A \circ f \circ A^{-1}, A \in \text{PSL}(2, \mathbb{C})$ .

**Def 5.4.4.** A point is called periodic if  $f^p(z) = z$  for some  $p \geq 1$  and its multiplier  $\lambda = (f^p)'(z)$ , then there are three cases:

- (1) when  $0 < |\lambda| < 1$ , we say  $z$  is attracting
- (2) when  $|\lambda| = 1$ , we say  $z$  is parabolic
- (3) when  $|\lambda| > 1$ , we say  $z$  is repelling.

A point is called critical point if  $f'(z) = 0$  and a point is called exceptional if  $O^-(z)$  is finite.

**Def 5.4.5.** Fatou set is  $F_f = \left\{ z \in \hat{\mathbb{C}} \mid \{f^n\} \text{ is a normal family in some neighborhood of } z \right\}$  and Julia set is  $J_f = \hat{\mathbb{C}} \setminus F_f$ .

**Thm 5.4.1.**  $J(f)$  is non-empty closed subset of  $\hat{\mathbb{C}}$ , both  $J(f)$  and  $F(f)$  are completely invariant under  $f$ .

*Proof.* Suppose  $J(f) = \emptyset$ , i.e.  $F(f) = \hat{\mathbb{C}}$ .

Then a subsequence  $\{f^{n_k}\}$  uniformly converges to some  $g : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ .

But  $\deg f^{n_k} \rightarrow \infty$ , contradiction! □

**Def 5.4.6.** For an attracting point  $z_0$ , we associate it with the basin of attraction

$$B(z_0) = \{z \in \hat{\mathbb{C}} \mid f^{pn}(z) \rightarrow z_0 \text{ as } n \rightarrow \infty\}.$$

**Lemma 5.4.1.**  $B(z_0) \subset F(f)$ .

**Lemma 5.4.2.** Repelling periodic points must be contained in  $J(f)$ .

*Proof.* Since  $f^p(z_0) = z_0$  and  $(f^p)'(z_0) > 1$ .

So  $(f^{pn})'(z_0) \rightarrow \infty$ . □

**Lemma 5.4.3.**  $J(f)$  is accumulation set of periodic points.

*Proof.* Let  $w \in J(f)$ , we may assume there exists  $v$  such that  $f(v) = w$  and  $f'(v) \neq 0$ .

In a neighborhood  $U$  of  $w$ , consider

$$h_k(z) = \frac{f^k(z) - z}{f^{-1}(z) - z}.$$

Then  $h_k$  is not a normal family in  $U$  since  $w \in J(f)$ .

So by Montel theorem, for some  $k$ ,  $h_k(z) = 0$  or  $h_k(z) = 1$  for  $z \in U$ .

Hence  $f^k(z) = z$  or  $f^k(z) = f^{-1}(z)$ , i.e.  $f^{k+1}(z) = z$ . □

**Exam 5.4.1.** Consider  $f_c = z^2 + c$ .

(1) If  $c = 0$ , then  $J(f_c) = \mathbb{S}^1$ .

(2) If  $0 < |c| < \frac{1}{4}$ , then  $J(f_c)$  is a quasicircle.

**Def 5.4.7.** A connected component  $U$  of  $F(f)$  is called a Fatou component. It is periodic if  $f^p(U) = U$  and it is pre-periodic if  $f^n(U)$  is periodic, otherwise it is called wandering.

**Thm 5.4.2.** A rational map of degree greater than one has no wandering domain.

**Def 5.4.8.** Suppose  $\mu$  is a Beltrami differential on  $\hat{\mathbb{C}}$  and  $f : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$  is a rational map, we say  $\mu$  is equivariant if  $(\mu \circ f) \frac{\bar{f}'}{f'} = \mu$  almost everywhere on  $\hat{\mathbb{C}}$ .

**Prop 5.4.1.** Suppose  $\phi : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$  such that  $\text{Belt}(\phi) = \mu$  is equivariant,  $g = \phi \circ f \circ \phi^{-1}$  is rational.

*Proof.*  $\text{Belt}(g \circ \phi) = \text{Belt}(\phi \circ f) = (\mu \circ f) \frac{\bar{f}'}{f'} = \mu$ .

So  $\text{Belt}(g) = 0$ , i.e.  $g$  is conformal. □

*Sketch of proof of theorem 5.4.2.* Suppose  $f : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$  has a wandering domain  $U$ , take  $\mu$  on  $U$ .

Then can be defined on  $O_{\text{grand}}(U)$  to be invariant under  $f$  and set  $\mu = 0$  outside  $O_{\text{grand}}(U)$ .

So we get  $\mu$  on  $\hat{\mathbb{C}}$  which is invariant under  $f$ .

Let  $f_t = \phi_t \circ f \circ \phi_t^{-1}$  where  $\text{Belt}(\phi_t) = t\mu$ ,  $V(z) = \frac{\partial}{\partial t} \phi_t(z)$  and  $\dot{f} = \frac{\partial}{\partial t} f_t$ .

By dimension count, there exists  $\mu$  such that  $\dot{f} \equiv 0$  and  $V \neq 0$  on  $\partial U \subset J(f)$ .

So  $[\mu] \neq 0$  in the tangent space of  $\text{Teich}(U)$ .

And since  $\dot{f}(z) = (V \circ f)(z) - f'(z)V(z)$ .

Therefore  $V \circ f = f' \cdot V$ .

Notice that  $J(f)$  is the closure of repelling periodic points.

So there exists  $z \in \partial U \subset J(f)$  such that  $f^n(z) = z$ ,  $V(z) \neq 0$  and  $(f^n)'(z) = \lambda$  with  $|\lambda| > 1$ .

Thus we obtain

$$\prod_{j=1}^n V(f^j(z)) = f'(f^{n-1}(z)) \cdots f'(f(z))f'(z) \prod_{j=0}^{n-1} V(f^j(z)) = (f^n)'(z) \prod_{j=1}^n V(f^j(z))$$

$$(\lambda - 1) \prod_{j=1}^n V(f^j(z)) = 0$$

But for every  $j$ ,

$$V(z) = V(f^n(z)) = f'(f^{n-1}(z)) \cdots f'(f^j(z))V(f^j(z)) \neq 0,$$

contradiction! □

# Appendix A

## Quasi-isometry and Mostow rigidity

### A.1 Boundary extension of quasi-isometry

**Lemma A.1.1.**  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$  is an  $(L, A)$ -quasi-isometry,  $\gamma$  is a geodesic ray and  $\gamma'$  is the geodesic ray within distance  $R$  from the quasigeodesic  $f(\gamma)$ . Let  $\pi : \mathbb{H}^n \rightarrow \gamma, \pi' : \mathbb{H}^n \rightarrow \gamma'$  be nearest-point restriction maps, then we have

$$d(f \circ \pi(x), \pi' \circ f(x)) \leq C,$$

for every  $x \in \mathbb{H}^n$  and some constant  $C$ .

This lemma is called the quasi-commute property, once we have done this proof, we can complete the proof of theorem 2.1.2.

**Thm A.1.1.**  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$  is a quasi-isometry then  $f$  extend continuous to a homeomorphism  $\partial f : \partial\mathbb{H}^n \rightarrow \partial\mathbb{H}^n$ .

*Proof.* By Morse lemma, let  $\gamma$  be the geodesic ray from 0 to  $a \in \partial\mathbb{H}^n$ ,  $f \circ \gamma$  be within a bounded distance  $R$  from  $\delta$  and define  $\partial f(\gamma(+\infty)) = \delta(+\infty)$ .

Consider a sequence  $\{x_i\}$  converging to  $a$ .

Then the boundary sphere  $\Sigma_i$  of  $H_i = \pi^{-1}(x_i)$  bound balls  $B_i \subset \mathbb{S}^{n-1}$  containing  $a$ .

So  $\{B_i\}$  forms a neighborhood basis of  $a$ .

And by the definition of quasi-isometry map,  $y_i = f(x_i)$  converges to  $\partial f(a)$ .

Let  $H'_i = (\pi')^{-1}(z_i)$  where  $d(z_i, \pi'(y_i)) = C$  and  $z_i, y_i, \partial f(a)$  are in order.

Then by lemma A.1.1,  $f(B_i) \subset B'_i$ .

So  $\partial f(B_i)$  forms a neighborhood basis of  $\partial f(a)$ , i.e.  $f$  is continuous.

Similarly, we can take balls  $B''_i \subset \mathbb{S}^{n-1}$  such that  $B''_i \subset \partial f(B_i)$ .

So  $f$  is open.

Moreover, suppose  $\partial f(a) = \partial f(b)$  and let  $\gamma_1, \gamma_2$  be the geodesic ray from 0 to  $a, b$  resp.

Then by Morse lemma  $f \circ \gamma_i$  are both within a bounded distance  $R$  from  $[f(0), \partial f(a)]$ .

So for any  $x \in \gamma_1$ , there exists  $y \in \gamma_2$  such that  $d(f(x), f(y)) \leq 2R$ .

Therefore  $d(x, y) \leq Ld(f(x), f(y)) + LA \leq 2LR + LA$  is bounded, i.e.  $\gamma_1, \gamma_2$  are asymptotic.

Thus  $a = b$ , i.e.  $f$  is injective.

Hence  $f$  is homeomorphic. □


*Proof of theorem 2.1.2.* WLOG, assume  $0, \infty \in \partial\mathbb{H}^n$  are the fixed points of  $\partial f$ .

Take  $x, y$  such that  $|x| = |y| = r$  and  $p \in \gamma$  where  $\gamma$  is the geodesic line connecting 0 and  $\infty$ .

Then by lemma A.1.1,

$$d(\pi(\partial f(x)), \pi(\partial f(y))) \leq d(f(\pi(x)), f(\pi(y))) + 2C \leq 2C$$

Notice that as the figure below,  $d(\pi(\partial f(x)), \pi(\partial f(y))) = |\ln |f(x)| - \ln |f(y)||$ .



So  $\frac{|f(x)-0|}{|f(y)-0|} \leq e^{2C}$ , i.e.  $\partial f$  is  $e^{2C}$ -quasiconformal.  $\square$

Now we only need to prove lemma A.1.1, we provide two different proofs. The second proof is more straightforward than the first one but I spent a week to complete the first proof before I suddenly conceived the second proof, whose most crucial step was inspired by the first proof. (Maybe this is because I am not so familiar with the hyperbolic geometry) So I decide to retain both.

For the first prove, we need some technique from Gromov. To simplify, we denote  $[x, y]$  as the geodesic segment from  $x$  to  $y$  below.

**Def A.1.1.** For three points  $u, v, w \in \mathbb{H}^n$ , the Gromov product of  $u, v$  at  $w$  is defined by

$$(u, v)_w = \frac{1}{2}(d(u, w) + d(v, w) - d(u, v)).$$

**Lemma A.1.2.** Let  $\gamma$  be a geodesic in  $\mathbb{H}^n$  and  $y \in \gamma$ , for a point  $x \in \mathbb{H}^n$ , take  $z \in \gamma$  such that  $d(x, z) = d(x, \gamma)$ , then prove that  $[x, z] + [z, y]$  is  $(\sqrt{2}, 0)$ -quasi-isometry.

*Proof.* Let  $\delta = [x, z] + [z, y]$  and  $\pi$  be the nearest point restriction map to  $[x, y]$ .

Reparametrize  $\delta$  by the length of image under  $\pi$ .

Since we have

$$\begin{aligned} \cosh^2 d(z, \pi(z)) &= \frac{\cosh d(x, z) \cosh d(y, z)}{\cosh d(x, \pi(z)) \cosh d(y, \pi(z))} \\ &= \frac{\cosh d(x, y)}{\cosh d(x, \pi(z)) \cosh d(y, \pi(z))} \\ &= 1 + \tanh d(x, \pi(z)) \tanh d(y, \pi(z)) \leq 2 \end{aligned}$$

So for any  $p \in \delta$ ,  $d(p, \pi(p)) \leq \cosh^{-1} \sqrt{2}$ .

Hence

$$|s - t| \leq d(\delta(s), \delta(t)) \leq \sqrt{2}d(\pi(\delta(s)), \pi(\delta(t))) = \sqrt{2}|s - t|$$

$\square$

**Lemma A.1.3.** Let  $\gamma : \mathbb{R} \rightarrow \mathbb{H}^n$  be an  $(L, A)$ -quasigeodesic and  $p, q, r$  are 3 points in order on  $\gamma$ , prove that there exists a constant  $K$  such that  $(p, r)_q \leq K$ .


*Proof.* Let  $\pi$  be the nearest point restriction map to  $[p, r]$ .

By Morse lemma,  $d(q, \pi(q)) \leq R$  for some  $R$  dependent only on  $L, A$ , so we have

$$\begin{aligned} (p, r)_q &= \frac{1}{2}(d(p, q) + d(r, q) - d(p, r)) \\ &= \frac{1}{2}(d(p, q) + d(r, q) - d(p, \pi(q)) - d(r, \pi(q))) \\ &\leq d(q, s) \leq R \end{aligned}$$

$\square$

proof of lemma A.1.1.



Let  $y = \pi(x)$ ,  $p = \pi'(f(x))$ ,  $q = \pi'(f(y))$  and  $r \in f \circ \gamma$  such that  $d(r, p) \leq R$ . By lemma A.1.2,  $[f(x), p] + [p, q]$  and  $[x, y] + [y, f^{-1}(r)]$  are quasi-isometry. So by lemma A.1.3,  $(f(x), q)_p \leq K_1$ ,  $(f(x), r)_{f(y)} \leq K_2$ .

Hence by Morse lemma,

$$\begin{aligned}
 d(f(y), p) &\leq d(f(y), q) + d(p, q) = d(f(y), q) + (q, f(x))_p + (p, f(x))_q \\
 &\leq R + K_1 + \frac{1}{2}(d(p, q) + d(f(x), q) - d(p, f(x))) \\
 &\leq R + K_1 + \frac{1}{2}(d(r, f(y)) + d(p, r) + d(f(y), q) \\
 &\quad + d(f(x), f(y)) + d(f(y), q) - d(r, f(x)) + d(p, r)) \\
 &= R + K_1 + d(f(y), q) + d(p, r) + (f(x), r)_{f(y)} \\
 &\leq K_1 + K_2 + 3R
 \end{aligned}$$

□

Notice that in lemma A.1.2, we use a counter-intuitive property of  $\mathbb{H}^n$ :

**Prop A.1.1.** Consider a geodesic right triangle  $\triangle$  in  $\mathbb{D}^n$ , let  $x$  be the right angle vertex and  $l$  be the hypotenuse, then  $d(x, l) \leq \operatorname{arccosh}\sqrt{2}$ .


*Proof.* We have given a proof by some hyperbolic trigonometry in the proof of lemma A.1.2, we now give a proof with some kind of geometric intuition.

WLOG, we assume  $x$  is at the origin.

Then legs of  $\triangle$  are on some radius of  $\mathbb{D}^n$ , denote them by  $l_1, l_2$ .

Let  $y, z$  be the boundary points of  $l_1, l_2$  resp.

So  $d(x, l) \leq d(x, [y, z]) = \operatorname{arccosh}\sqrt{2}$ .




□

*Alternative proof of lemma A.1.1.* Let  $a, b$  be the endpoints of  $\gamma$  and  $l_1 = [a, x], l_2 = [b, x]$ .

Then by Morse lemma,  $l'_1 = [f(a), f(x)], l'_2 = [f(b), f(y)]$  are within distance  $R$  from the quasigeodesic  $f(l_1), f(l_2)$  resp.

Let  $y = \pi(x)$ ,  $p = \pi'(f(x))$ ,  $q = \pi'(f(y))$  and  $r_i \in l'_i$  such that  $d(f(y), l_i) = d(f(y), r_i)$ , so

$$\begin{aligned}
 d(f(y), p) &\leq d(f(y), q) + d(p, q) \\
 &\leq R + d(q, \pi(r_1)) + d(q, \pi(r_2)) \\
 &\leq R + d(f(y), r_1) + d(f(y), r_2) \\
 &\leq R + d(f(y), f(l_1)) + R + d(f(y), f(l_2)) + R \\
 &\leq 3R + Ld(y, l_1) + A + Ld(y, l_2) + A \\
 &\leq 3R + 2L\text{Arccosh}\sqrt{2} + 2A
 \end{aligned}$$



□

**Prop A.1.2.** Suppose  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$  is a  $(L, A)$ -quasi-isometry, then  $f$  is quasi-dense: there exists a constant  $C$  such that for  $y \in \mathbb{H}^n$ , there is some  $x \in \mathbb{H}^n$  satisfying that

$$d(f(x), y) < C.$$

*Proof.* Take a geodesic line  $\gamma$  containing  $y$  which ends at  $a, b$ .

By theorem 2.1.2,  $p = \partial f^{-1}(a)$  and  $q = \partial f^{-1}(b)$  are well-defined.

Take the geodesic line  $\gamma'$  ending at  $p, q$ .

Then  $f \circ \gamma'$  is within a bounded distance  $R$  from  $\gamma$  where  $R$  only depends on  $L, A$ .

Let  $\pi$  be the nearest point restriction map for  $\gamma$ .

So  $\pi^{-1}(y) \cap f \circ \gamma'$  is nonempty, take  $p$  in it.

Hence  $p = f(x)$  for some  $x$  and

$$d(f(x), y) = d(p, y) = d(p, \pi(p)) \leq R.$$

□

*Remark A.1.1.* For general definition of quasi-isometry, we require the quasi-dense property. But since we can prove this property for  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$ , we did not required it when defining quasi-isometry  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$ .

**Prop A.1.3.**  $(L, A)$ -quasi-isometry map  $f : \mathbb{H}^n \rightarrow \mathbb{H}^n$  has a quasi-inverse: a  $(L, A_1)$ -quasi-isometry map  $g : \mathbb{H}^n \rightarrow \mathbb{H}^n$  such that

$$d(f \circ g(y), y) \leq B, d(g \circ f(x), x) \leq B.$$

*Proof.* Take  $g(y) \in \mathbb{H}^n$  such that  $d(f \circ g(y), y) \leq A$ , then

$$\begin{aligned}
 d(g(y_1), g(y_2)) &\leq Ld(f \circ g(y_1), f \circ g(y_2)) + LA \\
 &L(d(f \circ g(y_1), y_1) + d(f \circ g(y_2), y_2) + d(y_1, y_2)) + LA \\
 &\leq Ld(y_1, y_2) + 3LA
 \end{aligned}$$

Similarly,  $d(y_1, y_2) \leq Ld(g(y_1), g(y_2)) + 3A$ .

And since

$$d(g(f(x)), x) \leq Ld(f(g(f(x))), f(x)) + LA \leq 2LA.$$

So  $g$  is  $(L, A_1)$ -quasi-isometry for some  $A_1$  and it is quasi-inverse of  $f$ .  $\square$

**Prop A.1.4.** *Suppose  $f, g : \mathbb{H}^n \rightarrow \mathbb{H}^n$  are two  $(L, A)$ -quasi-isometry such that  $\partial f = \partial g$ , then there exists  $C > 0$  such that  $d(f(x), g(x)) \leq C$  for every  $x \in \mathbb{H}^n$ .*

*Proof.* Consider  $x \in \mathbb{H}^n$  and a geodesic line  $\gamma$  ending at  $a, b \in \partial\mathbb{H}^n$ .

Since  $\partial f(a) = \partial g(a), \partial f(b) = \partial g(b)$ .

So both  $f \circ \gamma, g \circ \gamma$  are within a bounded distance  $R$  from the geodesic line  $\gamma'$  ending at  $\partial f(a), \partial f(b)$ , where  $R$  only depends on  $L, A$ .

Therefore  $d(f(x), \gamma'), d(g(x), \gamma') \leq R$ .

Notice that  $\partial f$  is surjective.

In the other words, for every point  $p \in \partial\mathbb{H}^n$ , there exists a geodesic line  $\gamma_p$  ending at  $p$  such that  $\gamma_p$  intersects with both  $B_R(f(x))$  and  $B_R(g(x))$ .

Suppose  $U = B_R(f(x))$  and  $V = B_R(g(x))$  are disjoint.

let  $\delta$  be the geodesic segment from  $f(x)$  to  $g(x)$  and take a point  $q \in \delta \setminus (U \cup V)$ .

Consider the nearest point restriction map  $\pi$  for  $\delta$  and the hypersurface  $H = \pi^{-1}(q)$ .

Then  $H$  is totally geodesic and divides  $\overline{\mathbb{H}^n}$  into two components  $C_1, C_2$  where  $y_i \in C_i$ .

Moreover,  $U \subset C_1$  and  $V \subset C_2$  since  $\delta$  is perpendicular to  $H$ ,  $d(q, y_i) > R$ .

Take  $p \in H \cap \partial\mathbb{H}^n$ .

Since both  $\overline{C_1}, \overline{C_2}$  are convex.

So any geodesic line  $\gamma_p$  ending at  $p$  disjoints with one of  $U$  and  $V$ , contradiction!

Hence  $d(f(x), g(x)) \leq C$ .  $\square$

## A.2 Mostow rigidity theorem

**Thm A.2.1** (Mostow rigidity).  *$M, N$  are closed hyperbolic  $n$ -manifold with  $n \geq 3$ , then every homotopy equivalent  $f : M \rightarrow N$  must be homeomorphism to an isometry*

*Proof.* Let  $g : N \rightarrow M$  be the homotopy inverse of  $f$ .

By differential topology,  $f, g$  can be homotopy to some smooth maps, *i.e.* WLOG, we assume  $f, g$  are smooth.

And we can lift  $f, g$  to  $\tilde{f}, \tilde{g} : \mathbb{H}^n \rightarrow \mathbb{H}^n$ .

Then  $\tilde{f}, \tilde{g}$  are Lipschitz.

Moreover, there exists a smooth homotopy  $H : M \times I \rightarrow M$  such that  $g \circ f \stackrel{H}{\simeq} \text{Id}_M$ .

With some suitably  $\tilde{g}$ , we can lift  $H$  to  $\tilde{H}$  such that  $\tilde{g} \circ \tilde{f} \stackrel{\tilde{H}}{\simeq} \text{Id}_{\mathbb{H}^n}$ .

So  $\left| d\tilde{H} \right|$  is bounded, *i.e.* there exists constant  $C$  such that

$$d(\tilde{g} \circ \tilde{f}(x), x) \leq C.$$

Thus  $\tilde{f}$  is quasi-isometry with inverse  $\tilde{g}$ .

By theorem 2.1.2,  $\tilde{f}$  can be extended to quasiconformal  $\partial\tilde{f} : \partial\mathbb{H}^n \rightarrow \partial\mathbb{H}^n$ .

We claim that  $\partial\tilde{f}$  is actually conformal.

Once we have proved this, we can extend  $\partial\tilde{f}$  to a conformal map  $h : \mathbb{H}^n \rightarrow \mathbb{H}^n$ , *i.e.* it is an isometry under the hyperbolic metric.

Since for  $A \in \Gamma_X$ , there exists  $B \in \Gamma_Y$  such that  $\partial\tilde{f} \circ A = B \circ \partial\tilde{f}$ .

So  $h \circ A = B \circ h$  on  $\mathbb{H}^n$ , take

$$H_0 : \mathbb{H}^n \times I \rightarrow \mathbb{H}^n, (x, t) \mapsto (1-t)\tilde{f}(x) + th(x),$$

then  $\tilde{f} \stackrel{H_0}{\simeq} h$  and  $H_0 \circ A = B \circ \tilde{H}_0$ .

Hence  $H_0 : M \times I \rightarrow N$  is well-defined and  $f \stackrel{H_0}{\simeq} h$ .  $\square$

We now prove the claim in the proof of Mostow rigidity, it is a direct consequence of the theorem below.

**Thm A.2.2.** *Suppose  $X, Y$  are closed hyperbolic  $n$ -manifolds with  $n \geq 3$  and  $f : \partial\mathbb{H}^n \rightarrow \partial\mathbb{H}^n$  is a homeomorphism which is differentiable at  $x_0$  with nonzero derivative. If  $f \circ \gamma \circ f^{-1} \in \Gamma_Y$  for any  $\gamma \in \Gamma_X$ , then  $f$  is a Möbius transformation.*

*Proof.* WLOG, assume  $x_0 = 0$  and  $f(0) = 0$ .

Similar to proposition 2.2.2, we let  $A_n(x) = \frac{x}{n}$ , take  $y \in \mathbb{H}^n$  and a fundamental domain  $F$  of  $X$ , then we have  $\gamma_n \circ A_n(y) \in F$ ,  $\gamma_{n_i} \circ A_{n_i} \rightarrow \sigma_1$ ,  $\delta_n \circ f = f \circ \gamma_n$  and  $\delta_{n_i} \circ A_{n_i} \rightarrow \sigma_2$ .

Let  $f_n = A_n^{-1} \circ f \circ A_n$  converges to  $h : \partial\mathbb{H}^n \rightarrow \partial\mathbb{H}^n$  and  $J = f'(0)$ .

Then  $\sigma_2^{-1} \circ \delta \circ \sigma_2 \circ h = h \circ \sigma_1^{-1} \circ \gamma \circ \sigma_1$  and  $h(x) = Jx$  for  $x \in \partial\mathbb{H}^n$ .

Notice that

$$\begin{aligned} \sigma_2 \circ h \circ \sigma_1^{-1} &= \lim_{i \rightarrow \infty} \sigma_{n_i} \circ A_{n_i} \circ A_{n_i}^{-1} \circ f \circ A_{n_i} \circ A_{n_i}^{-1} \circ \gamma_{n_i}^{-1} \\ &= \lim_{i \rightarrow \infty} \sigma_{n_i} \circ f \circ \gamma_{n_i}^{-1} = f \end{aligned}$$

So we only need to prove that  $h$  is a Möbius transformation.

By Liouville theorem, this is equivalent to prove that  $h$  is an Euclidean similarity, which follows from the lemma below since  $\Gamma_X$  is cocompact.  $\square$

**Lemma A.2.1.** *Suppose  $\gamma \in \text{SO}^+(n, 1)$  such that  $\gamma(\infty) \neq \{0, \infty\}$  and  $A \in \text{GL}(n-1)$  which conjugates  $\gamma$  to  $A\gamma A^{-1} \in \text{SO}^+(n, 1)$ , then  $A$  is an Euclidean similarity.*

*Proof.* Suppose  $A$  is not an Euclidean similarity.

Since  $A\gamma(\infty) \neq 0$ , let  $P$  be a hyperplane in  $\mathbb{R}^{n-1}$  with  $0 \in P$ ,  $A\gamma(\infty) \notin P$ .

So  $\gamma A^{-1}(P)$  must be a sphere  $S \subset \mathbb{R}^{n-1}$  as  $\gamma$  is conformal.

But  $A(S)$  is an ellipsoid which is not a sphere.

Hence  $A\gamma A^{-1}$  is not conformal, contradiction!  $\square$